

# FZD5 Antibody

Purified Mouse Monoclonal Antibody

Catalog # AO1760a

## Product Information

---

<b>Application</b>	WB, FC, E
<b>Primary Accession</b>	<a href="#">Q13467</a>
<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	2D12
<b>Isotype</b>	IgG1
<b>Calculated MW</b>	64507
<b>Description</b>	Members of the 'frizzled' gene family encode 7-transmembrane domain proteins that are receptors for Wnt signaling proteins. The FZD5 protein is believed to be the receptor for the Wnt5A ligand.
<b>Immunogen</b>	Purified recombinant fragment of human FZD5 (AA:151-217) expressed in E. Coli.
<b>Formulation</b>	Purified antibody in PBS with 0.05% sodium azide

## Additional Information

---

<b>Gene ID</b>	7855
<b>Other Names</b>	Frizzled-5, Fz-5, hFz5, FzE5, FZD5, C2orf31
<b>Dilution</b>	WB~~1/500 - 1/2000 FC~~1/200 - 1/400 E~~1/10000
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	FZD5 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	FZD5
<b>Synonyms</b>	C2orf31
<b>Function</b>	Receptor for Wnt proteins (PubMed: <a href="#">10097073</a> , PubMed: <a href="#">20530549</a> , PubMed: <a href="#">26908622</a> , PubMed: <a href="#">9054360</a> ). Functions in the canonical Wnt/beta-catenin signaling pathway. In vitro activates WNT2, WNT10B, WNT5A, but not

WNT2B or WNT4 signaling (By similarity). In neurons, activation by WNT7A promotes formation of synapses (PubMed:[20530549](#)). May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues (Probable). Plays a role in yolk sac angiogenesis and in placental vascularization (By similarity). Plays a role in ocular development (PubMed:[26908622](#)).

## Cellular Location

Cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q8CHL0}. Golgi apparatus membrane {ECO:0000250|UniProtKB:Q9EQD0}; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q9EQD0}. Synapse {ECO:0000250|UniProtKB:Q8CHL0}. Perikaryon {ECO:0000250|UniProtKB:Q8CHL0}. Cell projection, dendrite {ECO:0000250|UniProtKB:Q8CHL0}. Cell projection, axon {ECO:0000250|UniProtKB:Q8CHL0}. Note=Localized at the plasma membrane and also found at the Golgi apparatus. {ECO:0000250|UniProtKB:Q9EQD0}

## References

1.J Biol Chem. 2009 Sep 25;284(39):26716-24.2.Int J Oncol. 2007 Mar;30(3):751-5.

## Images




Figure 1: Western blot analysis using FZD5 mAb against human FZD5 recombinant protein. (Expected MW is 32.5 kDa)




Figure 2: Western blot analysis using FZD5 mouse mAb against A549 (1), and PC-3 (2) cell lysate.




Figure 3: Flow cytometric analysis of HepG2 cells using FZD5 mouse mAb (green) and negative control (red).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.