

DCN Antibody

Purified Mouse Monoclonal Antibody

Catalog # AO1842a

Product Information

Application	WB, E
Primary Accession	P07585
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Clone Names	1G4C5
Isotype	IgG1
Calculated MW	39747
Description	The protein encoded by this gene is a small cellular or pericellular matrix proteoglycan that is closely related in structure to biglycan protein. The encoded protein and biglycan are thought to be the result of a gene duplication. This protein is a component of connective tissue, binds to type I collagen fibrils, and plays a role in matrix assembly. It contains one attached glycosaminoglycan chain. This protein is capable of suppressing the growth of various tumor cell lines. There are multiple alternatively spliced transcript variants known for this gene. This gene is a candidate gene for Marfan syndrome.
Immunogen	Purified recombinant fragment of human DCN (AA: 263-324) expressed in E. Coli.
Formulation	Purified antibody in PBS with 0.05% sodium azide

Additional Information

Gene ID	1634
Other Names	Decorin, Bone proteoglycan II, PG-S2, PG40, DCN, SLRR1B
Dilution	WB~~1/500 - 1/2000 E~~1/10000
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	DCN Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	DCN
-------------	-----

Synonyms	SLRR1B
Function	May affect the rate of fibrils formation.
Cellular Location	Secreted, extracellular space, extracellular matrix. Secreted
Tissue Location	Detected in placenta (at protein level) (PubMed:32337544). Detected in cerebrospinal fluid, fibroblasts and urine (at protein level) (PubMed:25326458, PubMed:36213313, PubMed:37453717).

Background

DNA (cytosine-5-)-methyltransferase 1 has a role in the establishment and regulation of tissue-specific patterns of methylated cytosine residues. Aberrant methylation patterns are associated with certain human tumors and developmental abnormalities. Two transcript variants encoding different isoforms have been found for this gene. ; ;

References

1. PLoS One. 2012;7(9):e45559. 2. Hum Reprod. 2012 Nov;27(11):3249-58.

Images




Figure 1: Western blot analysis using DCN mAb against human DCN recombinant protein. (Expected MW is 32.5 kDa)




Figure 2: Western blot analysis using DCN mAb against HEK293 (1) and DCN (AA: 263-324)-hIgGfC transfected HEK293 (2) cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.