

# CA9 Antibody

Purified Mouse Monoclonal Antibody

Catalog # AO1847a

## Product Information

---

<b>Application</b>	WB, IHC, E
<b>Primary Accession</b>	<a href="#">Q16790</a>
<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	7C4A5
<b>Isotype</b>	IgG1
<b>Calculated MW</b>	49698
<b>Description</b>	Carbonic anhydrases (CAs) are a large family of zinc metalloenzymes that catalyze the reversible hydration of carbon dioxide. They participate in a variety of biological processes, including respiration, calcification, acid-base balance, bone resorption, and the formation of aqueous humor, cerebrospinal fluid, saliva, and gastric acid. They show extensive diversity in tissue distribution and in their subcellular localization. CA IX is a transmembrane protein and the only tumor-associated carbonic anhydrase isoenzyme known. It is expressed in all clear-cell renal cell carcinoma, but is not detected in normal kidney or most other normal tissues. It may be involved in cell proliferation and transformation. This gene was mapped to 17q21.2 by fluorescence in situ hybridization, however, radiation hybrid mapping localized it to 9p13-p12.
<b>Immunogen</b>	Purified recombinant fragment of human CA9 (AA: 37-186) expressed in E. Coli.
<b>Formulation</b>	Purified antibody in PBS with 0.05% sodium azide

## Additional Information

---

<b>Gene ID</b>	768
<b>Other Names</b>	Carbonic anhydrase 9, 4.2.1.1, Carbonate dehydratase IX, Carbonic anhydrase IX, CA-IX, CAIX, Membrane antigen MN, P54/58N, Renal cell carcinoma-associated antigen G250, RCC-associated antigen G250, pMW1, CA9, G250, MN
<b>Dilution</b>	WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 E~~1/10000
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	CA9 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

<b>Name</b>	CA9
<b>Synonyms</b>	G250, MN
<b>Function</b>	Catalyzes the interconversion between carbon dioxide and water and the dissociated ions of carbonic acid (i.e. bicarbonate and hydrogen ions).
<b>Cellular Location</b>	Nucleus. Nucleus, nucleolus. Cell membrane; Single-pass type I membrane protein. Cell projection, microvillus membrane; Single-pass type I membrane protein. Note=Found on the surface microvilli and in the nucleus, particularly in nucleolus
<b>Tissue Location</b>	Expressed primarily in carcinoma cells lines. Expression is restricted to very few normal tissues and the most abundant expression is found in the epithelial cells of gastric mucosa

## Background

THY1 may play a role in cell-cell or cell-ligand interactions during synaptogenesis and other events in the brain ; ; ;

## References

1. Breast Cancer Res Treat. 2012 Nov;136(1):67-75. 2. Histol Histopathol. 2011 Oct;26(10):1279-86.

## Images




Figure 1: Western blot analysis using CA9 mAb against human CA9 recombinant protein. (Expected MW is 42 kDa)




Figure 2: Western blot analysis using CA9 mAb against HEK293 (1) and CA9 (AA: 37-186)-hIgGFc transfected HEK293 (2) cell lysate.

Figure 3: Western blot analysis using CA9 mouse mAb against A431 (1) and SW620 (2) cell lysate.





Figure 5: Immunohistochemical analysis of paraffin-embedded rectum cancer tissues using ZEB1 mouse mAb with DAB staining.




Figure 6: Immunohistochemical analysis of paraffin-embedded stomach tissues using CA9 mouse mAb with DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.