

# **ZEB1 Antibody**

Purified Mouse Monoclonal Antibody Catalog # AO1849a

#### **Product Information**

**Application** WB, IHC, FC, ICC, E

Primary Accession
Reactivity
Human
Host
Clonality
Monoclonal
Clone Names
Isotype
IgG1
Calculated MW
P37275
Human
Mouse
Clonal
2A8A6
IgG1
124074

**Description** This gene encodes a zinc finger transcription factor. The encoded protein

likely plays a role in transcriptional repression of interleukin 2. Mutations in this gene have been associated with posterior polymorphous corneal

dystrophy-3 and late-onset Fuchs endothelial corneal dystrophy. Alternatively spliced transcript variants encoding different isoforms have been described.

**Immunogen** Purified recombinant fragment of human ZEB1 (AA: 967-1108) expressed in E.

Coli.

**Formulation** Purified antibody in PBS with 0.05% sodium azide

#### **Additional Information**

**Gene ID** 6935

Other Names Zinc finger E-box-binding homeobox 1, NIL-2-A zinc finger protein, Negative

regulator of IL2, Transcription factor 8, TCF-8, ZEB1, AREB6, TCF8

**Dilution** WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 FC~~1/200 - 1/400 ICC~~N/A

E~~1/10000

**Storage** Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** ZEB1 Antibody is for research use only and not for use in diagnostic or

therapeutic procedures.

#### **Protein Information**

Name ZEB1 ( HGNC:11642)

**Function** Acts as a transcriptional repressor. Inhibits interleukin-2 (IL-2) gene

expression. Enhances or represses the promoter activity of the ATP1A1 gene depending on the quantity of cDNA and on the cell type. Represses E-cadherin promoter and induces an epithelial-mesenchymal transition (EMT) by recruiting SMARCA4/BRG1. Represses BCL6 transcription in the presence of the corepressor CTBP1. Positively regulates neuronal differentiation. Represses RCOR1 transcription activation during neurogenesis. Represses transcription by binding to the E box (5'-CANNTG-3'). In the absence of TGFB1, acts as a repressor of COL1A2 transcription via binding to the E-box in the upstream enhancer region (By similarity).

**Cellular Location** 

**Nucleus** 

**Tissue Location** 

Colocalizes with SMARCA4/BRG1 in E-cadherin- negative cells from established lines, and stroma of normal colon as well as in de-differentiated epithelial cells at the invasion front of colorectal carcinomas (at protein level). Expressed in heart and skeletal muscle, but not in liver, spleen, or pancreas

### **Background**

This gene encodes a zinc finger transcription factor. The encoded protein likely plays a role in transcriptional repression of interleukin 2. Mutations in this gene have been associated with posterior polymorphous corneal dystrophy-3 and late-onset Fuchs endothelial corneal dystrophy. Alternatively spliced transcript variants encoding different isoforms have been described.;;;

#### References

1. J Cancer Res Clin Oncol. 2012 Aug;138(8):1329-38. 2. Mol Cell Biochem. 2012 Jul;366(1-2):223-9.

## **Images**





Figure 1: Western blot analysis using ZEB1 mAb against human ZEB1 recombinant protein. (Expected MW is 41.7 kDa)

Figure 2: Western blot analysis using ZEB1 mAb against HEK293 (1) and ZEB1 (AA: 967-1108)-hIgGFc transfected HEK293 (2) cell lysate.

Figure 3: Immunofluorescence analysis of Hela cells using ZEB1 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.






Figure 4: Flow cytometric analysis of Hela cells using ZEB1 mouse mAb (green) and negative control (red).




Figure 5: Immunohistochemical analysis of paraffin-embedded cervical cancer tissues using ZEB1 mouse mAb with DAB staining.




Figure 6: Immunohistochemical analysis of paraffin-embedded rectum cancer tissues using ZEB1 mouse mAb with DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.