

# CDX2 Antibody

Purified Mouse Monoclonal Antibody

Catalog # AO1885a

## Product Information

---

<b>Application</b>	WB, FC, E
<b>Primary Accession</b>	<a href="#">Q99626</a>
<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	3G9B9
<b>Isotype</b>	IgG1
<b>Calculated MW</b>	33520
<b>Description</b>	This gene is a member of the caudal-related homeobox transcription factor gene family. The encoded protein is a major regulator of intestine-specific genes involved in cell growth and differentiation. This protein also plays a role in early embryonic development of the intestinal tract. Aberrant expression of this gene is associated with intestinal inflammation and tumorigenesis.
<b>Immunogen</b>	Purified recombinant fragment of human CDX2 (AA: 176-303) expressed in E. Coli.
<b>Formulation</b>	Purified antibody in PBS with 0.05% sodium azide

## Additional Information

---

<b>Gene ID</b>	1045
<b>Other Names</b>	Homeobox protein CDX-2, CDX-3, Caudal-type homeobox protein 2, CDX2, CDX3
<b>Dilution</b>	WB~~1/500 - 1/2000 FC~~1/200 - 1/400 E~~1/10000
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	CDX2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	CDX2
<b>Synonyms</b>	CDX3

<b>Function</b>	Transcription factor which regulates the transcription of multiple genes expressed in the intestinal epithelium (By similarity). Binds to the promoter of the intestinal sucrase-isomaltase SI and activates SI transcription (By similarity). Binds to the DNA sequence 5'-ATAAAACTTAT-3' in the promoter region of VDR and activates VDR transcription (By similarity). Binds to and activates transcription of LPH (By similarity). Activates transcription of CLDN2 and intestinal mucin MUC2 (By similarity). Binds to the 5'-AATTTTTTACAACACCT-3' DNA sequence in the promoter region of CA1 and activates CA1 transcription (By similarity). Important in broad range of functions from early differentiation to maintenance of the intestinal epithelial lining of both the small and large intestine. Binds preferentially to methylated DNA (PubMed: <a href="#">28473536</a> ).
<b>Cellular Location</b>	Nucleus {ECO:0000250 UniProtKB:P43241}.
<b>Tissue Location</b>	Detected in small intestine, colon and pancreas.

## Background

The protein encoded by this gene is a surface antigen that is preferentially expressed on monocytes/macrophages. It cooperates with other proteins to mediate the innate immune response to bacterial lipopolysaccharide. Alternative splicing results in multiple transcript variants encoding the same protein. ; ; ; ;

## References

1. Tumour Biol. 2012 Dec;33(6):2185-8. 2. Cancer Biol Ther. 2012 Oct;13(12):1152-7.

## Images




Figure 1: Western blot analysis using CDX2 mAb against human CDX2 (AA: 176-303) recombinant protein. (Expected MW is 40.1 kDa)





Figure 2: Western blot analysis using CDX2 mAb against HEK293 (1) and CDX2 (AA: 176-303)-hIgGFc transfected HEK293 (2) cell lysate.

Figure 3: Flow cytometric analysis of Hela cells using CDX2 mouse mAb (green) and negative control (red).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.