

# PRKAG1 Antibody

Purified Mouse Monoclonal Antibody

Catalog # AO1897a

## Product Information

---

<b>Application</b>	WB, IHC, FC, ICC, E
<b>Primary Accession</b>	<a href="#">P54619</a>
<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	7H3E9
<b>Isotype</b>	IgG1
<b>Calculated MW</b>	37579
<b>Description</b>	The protein encoded by this gene is a regulatory subunit of the AMP-activated protein kinase (AMPK). AMPK is a heterotrimer consisting of an alpha catalytic subunit, and non-catalytic beta and gamma subunits. AMPK is an important energy-sensing enzyme that monitors cellular energy status. In response to cellular metabolic stresses, AMPK is activated, and thus phosphorylates and inactivates acetyl-CoA carboxylase (ACC) and beta-hydroxy beta-methylglutaryl-CoA reductase (HMGCR), key enzymes involved in regulating de novo biosynthesis of fatty acid and cholesterol. This subunit is one of the gamma regulatory subunits of AMPK. Alternatively spliced transcript variants encoding distinct isoforms have been observed.
<b>Immunogen</b>	Purified recombinant fragment of human PRKAG1 (AA: 230-331) expressed in E. Coli.
<b>Formulation</b>	Purified antibody in PBS with 0.05% sodium azide.

## Additional Information

---

<b>Gene ID</b>	5571
<b>Other Names</b>	5'-AMP-activated protein kinase subunit gamma-1, AMPK gamma1, AMPK subunit gamma-1, AMPKg, PRKAG1
<b>Dilution</b>	WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 FC~~1/200 - 1/400 ICC~~N/A E~~1/10000
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	PRKAG1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

Name	PRKAG1
Function	AMP/ATP-binding subunit of AMP-activated protein kinase (AMPK), an energy sensor protein kinase that plays a key role in regulating cellular energy metabolism (PubMed: <a href="#">21680840</a> , PubMed: <a href="#">24563466</a> ). In response to reduction of intracellular ATP levels, AMPK activates energy-producing pathways and inhibits energy-consuming processes: inhibits protein, carbohydrate and lipid biosynthesis, as well as cell growth and proliferation (PubMed: <a href="#">21680840</a> , PubMed: <a href="#">24563466</a> ). AMPK acts via direct phosphorylation of metabolic enzymes, and by longer-term effects via phosphorylation of transcription regulators (PubMed: <a href="#">21680840</a> , PubMed: <a href="#">24563466</a> ). Also acts as a regulator of cellular polarity by remodeling the actin cytoskeleton; probably by indirectly activating myosin (PubMed: <a href="#">21680840</a> , PubMed: <a href="#">24563466</a> ). Gamma non-catalytic subunit mediates binding to AMP, ADP and ATP, leading to activate or inhibit AMPK: AMP-binding results in allosteric activation of alpha catalytic subunit (PRKAA1 or PRKAA2) both by inducing phosphorylation and preventing dephosphorylation of catalytic subunits (PubMed: <a href="#">21680840</a> , PubMed: <a href="#">24563466</a> ). ADP also stimulates phosphorylation, without stimulating already phosphorylated catalytic subunit (PubMed: <a href="#">21680840</a> , PubMed: <a href="#">24563466</a> ). ATP promotes dephosphorylation of catalytic subunit, rendering the AMPK enzyme inactive (PubMed: <a href="#">21680840</a> , PubMed: <a href="#">24563466</a> ).

## Background

This gene encodes a serum protein found in association with the major histocompatibility complex (MHC) class I heavy chain on the surface of nearly all nucleated cells. The protein has a predominantly beta-pleated sheet structure that can form amyloid fibrils in some pathological conditions. A mutation in this gene has been shown to result in hypercatabolic hypoproteinemia. ;

## References

1. Circ Res. 2012 Aug 31;111(6):800-14.
2. Circ Res. 2012 Apr 27;110(9):1192-201.

## Images




Figure 1: Western blot analysis using PRKAG1 mAb against human PRKAG1 (AA: 230-331) recombinant protein. (Expected MW is 37.4 kDa)

Figure 2: Western blot analysis using PRKAG1 mAb against HEK293 (1) and PRKAG1 (AA: 230-331)-hIgGFc transfected HEK293 (2) cell lysate.




Figure 3: Immunofluorescence analysis of HepG2 cells using PRKAG1 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Secondary antibody from Fisher (Cat#: 35503)




Figure 4: Flow cytometric analysis of HepG2 cells using PRKAG1 mouse mAb (green) and negative control (red).




Figure 5: Immunohistochemical analysis of paraffin-embedded prostate cancer tissues using PRKAG1 mouse mAb with DAB staining.





Figure 6: Immunohistochemical analysis of paraffin-embedded kidney tissues using PRKAG1 mouse mAb with DAB staining.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.