

# NBN Antibody

Purified Mouse Monoclonal Antibody

Catalog # AO1911a

## Product Information

---

<b>Application</b>	WB, IHC, FC, ICC, E
<b>Primary Accession</b>	<a href="#">O60934</a>
<b>Reactivity</b>	Human, Rat
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	7E4A2
<b>Isotype</b>	IgG2a
<b>Calculated MW</b>	84959
<b>Description</b>	Mutations in this gene are associated with Nijmegen breakage syndrome, an autosomal recessive chromosomal instability syndrome characterized by microcephaly, growth retardation, immunodeficiency, and cancer predisposition. The encoded protein is a member of the MRE11/RAD50 double-strand break repair complex which consists of 5 proteins. This gene product is thought to be involved in DNA double-strand break repair and DNA damage-induced checkpoint activation.
<b>Immunogen</b>	Purified recombinant fragment of human NBN (AA: 467-615) expressed in E. Coli.
<b>Formulation</b>	Purified antibody in PBS with 0.05% sodium azide.

## Additional Information

---

<b>Gene ID</b>	4683
<b>Other Names</b>	Nibrin, Cell cycle regulatory protein p95, Nijmegen breakage syndrome protein 1, NBN, NBS, NBS1, P95
<b>Dilution</b>	WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 FC~~1/200 - 1/400 ICC~~N/A E~~1/10000
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	NBN Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	NBN ( <a href="#">HGNC:7652</a> )
-------------	-----------------------------------

## Function

Component of the MRN complex, which plays a central role in double-strand break (DSB) repair, DNA recombination, maintenance of telomere integrity and meiosis (PubMed:[10888888](#), PubMed:[15616588](#), PubMed:[18411307](#), PubMed:[18583988](#), PubMed:[18678890](#), PubMed:[19759395](#), PubMed:[23115235](#), PubMed:[28216226](#), PubMed:[28867292](#), PubMed:[9705271](#)). The MRN complex is involved in the repair of DNA double-strand breaks (DSBs) via homologous recombination (HR), an error-free mechanism which primarily occurs during S and G2 phases (PubMed:[19759395](#), PubMed:[28867292](#), PubMed:[9705271](#)). The complex (1) mediates the end resection of damaged DNA, which generates proper single-stranded DNA, a key initial steps in HR, and is (2) required for the recruitment of other repair factors and efficient activation of ATM and ATR upon DNA damage (PubMed:[19759395](#), PubMed:[9705271](#)). The MRN complex possesses single-strand endonuclease activity and double-strand-specific 3'-5' exonuclease activity, which are provided by MRE11, to initiate end resection, which is required for single-strand invasion and recombination (PubMed:[19759395](#), PubMed:[28867292](#), PubMed:[9705271](#)). Within the MRN complex, NBN acts as a protein-protein adapter, which specifically recognizes and binds phosphorylated proteins, promoting their recruitment to DNA damage sites (PubMed:[12419185](#), PubMed:[15616588](#), PubMed:[18411307](#), PubMed:[18582474](#), PubMed:[18583988](#), PubMed:[18678890](#), PubMed:[19759395](#), PubMed:[19804756](#), PubMed:[23762398](#), PubMed:[24534091](#), PubMed:[27814491](#), PubMed:[27889449](#), PubMed:[33836577](#)). Recruits MRE11 and RAD50 components of the MRN complex to DSBs in response to DNA damage (PubMed:[12419185](#), PubMed:[18411307](#), PubMed:[18583988](#), PubMed:[18678890](#), PubMed:[24534091](#), PubMed:[26438602](#)). Promotes the recruitment of PI3/PI4-kinase family members ATM, ATR, and probably DNA-PKcs to the DNA damage sites, activating their functions (PubMed:[15064416](#), PubMed:[15616588](#), PubMed:[15790808](#), PubMed:[16622404](#), PubMed:[22464731](#), PubMed:[30952868](#), PubMed:[35076389](#)). Mediates the recruitment of phosphorylated RBBP8/CtIP to DSBs, leading to cooperation between the MRN complex and RBBP8/CtIP to initiate end resection (PubMed:[19759395](#), PubMed:[27814491](#), PubMed:[27889449](#), PubMed:[33836577](#)). RBBP8/CtIP specifically promotes the endonuclease activity of the MRN complex to clear DNA ends containing protein adducts (PubMed:[27814491](#), PubMed:[27889449](#), PubMed:[30787182](#), PubMed:[33836577](#)). The MRN complex is also required for the processing of R-loops (PubMed:[31537797](#)). NBN also functions in telomere length maintenance via its interaction with TERF2: interaction with TERF2 during G1 phase preventing recruitment of DCLRE1B/Apollo to telomeres (PubMed:[10888888](#), PubMed:[28216226](#)). NBN also promotes DNA repair choice at dysfunctional telomeres: NBN phosphorylation by CDK2 promotes non-homologous end joining repair at telomeres, while unphosphorylated NBN promotes microhomology-mediated end-joining (MMEJ) repair (PubMed:[28216226](#)). Enhances AKT1 phosphorylation possibly by association with the mTORC2 complex (PubMed:[23762398](#)).

## Cellular Location

Nucleus. Chromosome. Nucleus, PML body. Chromosome, telomere  
Note=Localizes to discrete nuclear foci after treatment with genotoxic agents (PubMed:[10783165](#), PubMed:[26215093](#), PubMed:[26438602](#)). Localizes to DNA double-strand breaks (DSBs); recruited to DNA damage sites via association with phosphorylated proteins, such as phosphorylated H2AX, phosphorylated MDC1 and phosphorylated RAD17 (PubMed:[12419185](#), PubMed:[18411307](#), PubMed:[18582474](#), PubMed:[18583988](#), PubMed:[18678890](#), PubMed:[19338747](#), PubMed:[23115235](#), PubMed:[24534091](#), PubMed:[26438602](#)) Acetylation of 'Lys-5' of histone H2AX (H2AXK5ac) promotes NBN/NBS1 assembly at the sites of DNA damage (PubMed:[26438602](#))

## Tissue Location

Ubiquitous (PubMed:9590180). Expressed at high levels in testis (PubMed:9590180).

## Background

This gene is a member of the solute carrier family 2 (facilitated glucose transporter) family and encodes a protein that functions as an insulin-regulated facilitative glucose transporter. In the absence of insulin, this integral membrane protein is sequestered within the cells of muscle and adipose tissue. Within minutes of insulin stimulation, the protein moves to the cell surface and begins to transport glucose across the cell membrane. Mutations in this gene have been associated with noninsulin-dependent diabetes mellitus (NIDDM). ; ; ;

## References

1. Fam Cancer. 2012 Dec;11(4):595-600. 2. Mol Carcinog. 2011 Sep;50(9):689-96.

## Images




Figure 1: Western blot analysis using NBN mAb against human NBN (AA: 467-615) recombinant protein. (Expected MW is 44.3 kDa)




Figure 2: Western blot analysis using NBN mAb against HEK293 (1) and NBN (AA: 467-615)-hIgGFc transfected HEK293 (2) cell lysate.




Figure 3: Western blot analysis using NBN mouse mAb against A549 (1), Jurkat (2) and PC-12 (3) cell lysate.

Figure 4: Immunofluorescence analysis of Hela cells using NBN mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin. Secondary antibody from Fisher (Cat#: 35503)




Figure 5: Flow cytometric analysis of HeLa cells using NBN mouse mAb (green) and negative control (red).





Figure 6: Immunohistochemical analysis of paraffin-embedded cervical cancer tissues using NBN mouse mAb with DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.