

PTPN1 Antibody

Purified Mouse Monoclonal Antibody Catalog # AO1991a

Product Information

WB, FC, E Application **Primary Accession** P18031 Reactivity Human Host Mouse Clonality Monoclonal **Clone Names** 4F11B9 Isotype IgG1 **Calculated MW** 49967

Description The protein encoded by this gene is the founding member of the protein

tyrosine phosphatase (PTP) family, which was isolated and identified based on its enzymatic activity and amino acid sequence. PTPs catalyze the hydrolysis of the phosphate monoesters specifically on tyrosine residues. Members of the PTP family share a highly conserved catalytic motif, which is essential for the catalytic activity. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP has been shown to act as a

negative regulator of insulin signaling by dephosphorylating the

phosphotryosine residues of insulin receptor kinase. This PTP was also reported to dephosphorylate epidermal growth factor receptor kinase, as well as JAK2 and TYK2 kinases, which implicated the role of this PTP in cell growth control, and cell response to interferon stimulation. Two transcript variants

encoding different isoforms have been found for this gene.

Immunogen Purified recombinant fragment of human PTPN1 (AA: 40-246) expressed in E.

Coli.

Formulation Purified antibody in PBS with 0.05% sodium azide.

Additional Information

Gene ID 5770

Other Names Tyrosine-protein phosphatase non-receptor type 1, 3.1.3.48, Protein-tyrosine

phosphatase 1B, PTP-1B, PTPN1, PTP1B

Dilution WB~~1/500 - 1/2000 FC~~1/200 - 1/400 E~~1/10000

Storage Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions PTPN1 Antibody is for research use only and not for use in diagnostic or

therapeutic procedures.

Protein Information

Name PTPN1

Synonyms PTP1B

Function Tyrosine-protein phosphatase which acts as a regulator of endoplasmic

reticulum unfolded protein response. Mediates dephosphorylation of EIF2AK3/PERK; inactivating the protein kinase activity of EIF2AK3/PERK. May play an important role in CKII- and p60c- src-induced signal transduction cascades. May regulate the EFNA5-EPHA3 signaling pathway which modulates cell reorganization and cell-cell repulsion. May also regulate the hepatocyte growth factor receptor signaling pathway through dephosphorylation of MET.

Cellular Location Endoplasmic reticulum membrane; Peripheral membrane protein;

Cytoplasmic side Note=Interacts with EPHA3 at the cell membrane

Tissue Location Expressed in keratinocytes (at protein level).

Background

The protein encoded by this gene is a transmembrane glycoprotein that is a member of the protein kinase superfamily. This protein is a receptor for members of the epidermal growth factor family. EGFR is a cell surface protein that binds to epidermal growth factor. Binding of the protein to a ligand induces receptor dimerization and tyrosine autophosphorylation and leads to cell proliferation. Mutations in this gene are associated with lung cancer. Multiple alternatively spliced transcript variants that encode different protein isoforms have been found for this gene.;

References

1. Med Oncol. 2012 Jun;29(2):948-56. 2. Cell Biol Int. 2010 Jul;34(7):747-53.

Images




Figure 1: Western blot analysis using PTPN1 mAb against human PTPN1 (AA: 40-246) recombinant protein. (Expected MW is 50 kDa)

Figure 2: Western blot analysis using PTPN1 mAb against HEK293 (1) and PTPN1 (AA: 40-246)-hIgGFc transfected HEK293 (2) cell lysate.




Figure 3: Western blot analysis using PTPN1 mouse mAb against A431 cell lysate.




Figure 4: Flow cytometric analysis of A431 cells using PTPN1 mouse mAb (green) and negative control (red).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.