

CD6 Antibody

Purified Mouse Monoclonal Antibody

Catalog # AO2011a

Product Information

Application	WB, IHC, FC, ICC, E
Primary Accession	P30203
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Clone Names	5A10A2
Isotype	IgG2b
Calculated MW	71801
Description	This gene encodes a protein found on the outer membrane of T-lymphocytes as well as some other immune cells. The encoded protein contains three scavenger receptor cysteine-rich (SRCR) domains and a binding site for an activated leukocyte cell adhesion molecule. The gene product is important for continuation of T cell activation. This gene may be associated with susceptibility to multiple sclerosis (PMID: 19525953, 21849685). Multiple transcript variants encoding different isoforms have been found for this gene.
Immunogen	Purified recombinant fragment of human CD6 (AA: 18-199) expressed in E. Coli.
Formulation	Purified antibody in PBS with 0.05% sodium azide

Additional Information

Gene ID	923
Other Names	T-cell differentiation antigen CD6, T12, TP120, CD6, CD6
Dilution	WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 FC~~1/200 - 1/400 ICC~~N/A E~~1/10000
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	CD6 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CD6 (HGNC:1691)
-------------	-----------------------------------

Function	Cell adhesion molecule that mediates cell-cell contacts and regulates T-cell responses via its interaction with ALCAM/CD166 (PubMed: 15048703 , PubMed: 15294938 , PubMed: 16352806 , PubMed: 16914752 , PubMed: 24584089 , PubMed: 24945728). Contributes to signaling cascades triggered by activation of the TCR/CD3 complex (PubMed: 24584089). Functions as a costimulatory molecule; promotes T-cell activation and proliferation (PubMed: 15294938 , PubMed: 16352806 , PubMed: 16914752). Contributes to the formation and maturation of the immunological synapse (PubMed: 15294938 , PubMed: 16352806). Functions as a calcium- dependent pattern receptor that binds and aggregates both Gram-positive and Gram-negative bacteria. Binds both lipopolysaccharide (LPS) from Gram-negative bacteria and lipoteichoic acid from Gram-positive bacteria (PubMed: 17601777). LPS binding leads to the activation of signaling cascades and down-stream MAP kinases (PubMed: 17601777). Mediates activation of the inflammatory response and the secretion of pro-inflammatory cytokines in response to LPS (PubMed: 17601777).
Cellular Location	Cell membrane; Single-pass type I membrane protein. Note=Detected at the immunological synapse, i.e, at the contact zone between antigen-presenting dendritic cells and T-cells (PubMed:15294938, PubMed:16352806). Colocalizes with the TCR/CD3 complex at the immunological synapse (PubMed:15294938)
Tissue Location	Detected on thymocytes (PubMed:15294938). Detected on peripheral blood T-cells (PubMed:15048703, PubMed:16352806) Detected on natural killer (NK) cells (PubMed:16352806). Soluble CD6 is detected in blood serum (at protein level) (PubMed:17601777). Detected in spleen, thymus, appendix, lymph node and peripheral blood leukocytes (PubMed:9013954). Expressed by thymocytes, mature T-cells, a subset of B-cells known as B-1 cells, and by some cells in the brain

References

J Autoimmun. 2010 Dec;35(4):336-41.Proc Natl Acad Sci U S A. 2007 Jul 10;104(28):11724-9.

Images

