

# LHCGR Antibody

Purified Mouse Monoclonal Antibody Catalog # AO2014a

#### **Product Information**

**Application** WB, FC, ICC, E **Primary Accession** P22888 Reactivity Human Host Mouse Monoclonal Clonality **Clone Names** 8G9A2 Isotype IgG1 78643 **Calculated MW** 

**Description** This gene encodes the receptor for both luteinizing hormone and

choriogonadotropin. This receptor belongs to the G-protein coupled receptor 1 family, and its activity is mediated by G proteins which activate adenylate cyclase. Mutations in this gene result in disorders of male secondary sexual character development, including familial male precocious puberty, also known as testotoxicosis, hypogonadotropic hypogonadism, Leydig cell adenoma with precocious puberty, and male pseudohermaphtoditism with

Leydig cell hypoplasia.

**Immunogen** LHR-29 from ATCC

**Formulation** Purified antibody in PBS with 0.05% sodium azide

### **Additional Information**

**Gene ID** 3973

Other Names Lutropin-choriogonadotropic hormone receptor, LH/CG-R, Luteinizing

hormone receptor, LHR, LSH-R, LHCGR, LCGR, LGR2, LHRHR

**Dilution** WB~~1/500 - 1/2000 FC~~1/200 - 1/400 ICC~~N/A E~~1/10000

**Storage** Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** LHCGR Antibody is for research use only and not for use in diagnostic or

therapeutic procedures.

#### **Protein Information**

Name LHCGR

**Synonyms** LCGR, LGR2, LHRHR

**Function** Receptor for lutropin-choriogonadotropic hormone (PubMed: <u>11847099</u>). The

activity of this receptor is mediated by G proteins which activate adenylate

cyclase (PubMed:<u>11847099</u>).

**Cellular Location** Cell membrane; Multi-pass membrane protein

**Tissue Location** Gonadal and thyroid cells.

## References

Cancer Res. 57: 857-864, 1997 Reprod Biol Endocrinol. 2011 May 15;9:64.

## **Images**

