

DDX39B Antibody

Purified Mouse Monoclonal Antibody
Catalog # AO2125a

Product Information

Application	WB, IHC, E
Primary Accession	Q13838
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Clone Names	3A2B2
Isotype	IgG1
Calculated MW	48991 Da
Description	This gene encodes a member of the DEAD box family of RNA-dependent ATPases that mediate ATP hydrolysis during pre-mRNA splicing. The encoded protein is an essential splicing factor required for association of U2 small nuclear ribonucleoprotein with pre-mRNA, and it also plays an important role in mRNA export from the nucleus to the cytoplasm. This gene belongs to a cluster of genes localized in the vicinity of the genes encoding tumor necrosis factor alpha and tumor necrosis factor beta. These genes are all within the human major histocompatibility complex class III region. Mutations in this gene may be associated with rheumatoid arthritis. Alternative splicing results in multiple transcript variants. Related pseudogenes have been identified on both chromosomes 6 and 11. Read-through transcription also occurs between this gene and the upstream ATP6V1G2 (ATPase, H+ transporting, lysosomal 13kDa, V1 subunit G2) gene.
Immunogen	Purified recombinant fragment of human DDX39B (AA: 1-250) expressed in E. Coli.
Formulation	Purified antibody in PBS with 0.05% sodium azide

Additional Information

Other Names	Spliceosome RNA helicase DDX39B, 3.6.4.13, 56 kDa U2AF65-associated protein, ATP-dependent RNA helicase p47, DEAD box protein UAP56, HLA-B-associated transcript 1 protein, DDX39B, BAT1, UAP56
Dilution	WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 E~~1/10000
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	DDX39B Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

References

1. J Virol. 2011 Sep;85(17):8646-55. 2. Biochem Biophys Res Commun. 2010 Feb 26;393(1):106-10.

Images

