

MMP14 Antibody

Purified Mouse Monoclonal Antibody

Catalog # AO2127a

Product Information

Application	WB, IHC, FC, E
Primary Accession	P50281
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Clone Names	6A11D1
Isotype	IgG1
Calculated MW	65894
Description	Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. However, the protein encoded by this gene is a member of the membrane-type MMP (MT-MMP) subfamily; each member of this subfamily contains a potential transmembrane domain suggesting that these proteins are expressed at the cell surface rather than secreted. This protein activates MMP2 protein, and this activity may be involved in tumor invasion.
Immunogen	Purified recombinant fragment of human MMP14 (AA: 112-246) expressed in E. Coli.
Formulation	Purified antibody in PBS with 0.05% sodium azide

Additional Information

Gene ID	4323
Other Names	Matrix metalloproteinase-14, MMP-14, 3.4.24.80, MMP-X1, Membrane-type matrix metalloproteinase 1, MT-MMP 1, MTMMP1, Membrane-type-1 matrix metalloproteinase, MT1-MMP, MT1MMP, MMP14
Dilution	WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 FC~~1/200 - 1/400 E~~1/10000
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	MMP14 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	MMP14
Function	Endopeptidase that degrades various components of the extracellular matrix such as collagen (PubMed: 8015608). Essential for pericellular collagenolysis and modeling of skeletal and extraskeletal connective tissues during development (By similarity). Activates progelatinase A/MMP2, thereby acting as a positive regulator of cell growth and migration (PubMed: 22065321 , PubMed: 8015608). Involved in the formation of the fibrovascular tissues in association with pro-MMP2 (PubMed: 12714657 , PubMed: 22065321). May be involved in actin cytoskeleton reorganization by cleaving PTK7 (PubMed: 20837484). Acts as a regulator of Notch signaling by mediating cleavage and inhibition of DLL1 (PubMed: 21572390). Cleaves ADGRB1 to release vasculostatin-40 which inhibits angiogenesis (PubMed: 22330140). Acts as a negative regulator of the GDF15-GFRAL aversive response by mediating cleavage and inactivation of GFRAL (PubMed: 35177851).
Cellular Location	Cell membrane; Single-pass type I membrane protein. Melanosome. Cytoplasm Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed: 17081065). Forms a complex with BST2 and localizes to the cytoplasm (PubMed: 17081065)
Tissue Location	Expressed in stromal cells of colon, breast, and head and neck. Expressed in lung tumors.

References

1.Cancer Med. 2013 Aug;2(4):457-67.2.BMC Cancer. 2013 Feb 10;13:74.

Images

