

# RAD23A Antibody

Purified Mouse Monoclonal Antibody  
Catalog # AO2165a

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">P54725</a>
<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	1E4D6
<b>Isotype</b>	IgG2b
<b>Calculated MW</b>	39609
<b>Description</b>	The protein encoded by this gene is one of two human homologs of <i>Saccharomyces cerevisiae</i> Rad23, a protein involved in nucleotide excision repair. Proteins in this family have a modular domain structure consisting of an ubiquitin-like domain (Ubl), ubiquitin-associated domain 1 (Uba1), XPC-binding domain and Uba2. The protein encoded by this gene plays an important role in nucleotide excision repair and also in delivery of polyubiquitinated proteins to the proteasome. Alternative splicing results in multiple transcript variants encoding multiple isoforms.
<b>Immunogen</b>	Purified recombinant fragment of human RAD23A (AA: 1-363) expressed in <i>E. Coli</i> .
<b>Formulation</b>	Purified antibody in PBS with 0.05% sodium azide

## Additional Information

---

<b>Gene ID</b>	5886
<b>Other Names</b>	UV excision repair protein RAD23 homolog A, HR23A, hHR23A, RAD23A
<b>Dilution</b>	WB~1/500 - 1/2000 E~1/10000
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	RAD23A Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name RAD23A ( [HGNC:9812](#) )

## Function

Multifunctional protein that participates in histone H4K20 demethylation, DNA repair, ubiquitin-dependent protein degradation, transcriptional regulation, and viral replication (PubMed:[12643283](#), PubMed:[14621999](#), PubMed:[15321727](#), PubMed:[20614012](#), PubMed:[32209475](#), PubMed:[9372924](#)). Specifically demethylates mono-, di- and trimethylated 'Lys-20' of histone H4 (H4K20me1, H4K20me2, H4K20me3, respectively) into unmethylated forms. Activates the transcription of coding genes by demethylating H4K20me1 and the transcription of repetitive elements by demethylating H4K20me3 (PubMed:[32209475](#)). Involved in modulation of proteasomal degradation. Binds to 'Lys-48'-linked polyubiquitin chains in a length-dependent manner and with a lower affinity to 'Lys-63'- linked polyubiquitin chains. Proposed to be capable to bind simultaneously to the 26S proteasome and to polyubiquitinated substrates and to deliver ubiquitinated proteins to the proteasome (PubMed:[12643283](#), PubMed:[14621999](#), PubMed:[15321727](#)). Involved in nucleotide excision repair, is considered functionally equivalent to RAD23B in global genome nucleotide excision repair (GG-NER) through its association with XPC. In vitro, the XPC-RAD23A complex demonstrates NER activity (PubMed:[9372924](#)). Can stabilize XPC (By similarity).

## Cellular Location

Nucleus.

## References

1. Biochem Biophys Res Commun. 2013 Feb 22;431(4):686-92. 2. Exp Cell Res. 2001 Jul 15;267(2):243-57.

## Images

