

ULK2 Antibody

Purified Mouse Monoclonal Antibody Catalog # AO2209a

Product Information

Application WB, FC, E **Primary Accession** QSIYT8

Reactivity Human, Mouse, Rat

Host Mouse
Clonality Monoclonal
Clone Names 2H4B2
Isotype IgG1
Calculated MW 112694

Description This gene encodes a protein that is similar to a serine/threonine kinase in C.

elegans which is involved in axonal elongation. The structure of this protein is similar to the C. elegans protein in that both proteins have an N-terminal kinase domain, a central proline/serine rich (PS) domain, and a C-terminal (C) domain. The gene is located within the Smith-Magenis syndrome region on chromosome 17. Alternatively spliced transcript variants encoding the same

protein have been identified.

Immunogen Purified recombinant fragment of human ULK2 (AA: 1-155) expressed in E.

Coli.

Formulation Purified antibody in PBS with 0.05% sodium azide

Additional Information

Gene ID 9706

Other Names Serine/threonine-protein kinase ULK2, 2.7.11.1, Unc-51-like kinase 2, ULK2,

KIAA0623

Dilution WB~~1/500 - 1/2000 FC~~1/200 - 1/400 E~~1/10000

Storage Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions ULK2 Antibody is for research use only and not for use in diagnostic or

therapeutic procedures.

Protein Information

Name ULK2

Synonyms

KIAA0623

Function

Serine/threonine-protein kinase involved in autophagy in response to starvation. Acts upstream of phosphatidylinositol 3-kinase PIK3C3 to regulate the formation of autophagophores, the precursors of autophagosomes. Part of regulatory feedback loops in autophagy: acts both as a downstream effector and a negative regulator of mammalian target of rapamycin complex 1 (mTORC1) via interaction with RPTOR. Activated via phosphorylation by AMPK, also acts as a negative regulator of AMPK through phosphorylation of the AMPK subunits PRKAA1, PRKAB2 and PRKAG1. May phosphorylate ATG13/KIAA0652, FRS2, FRS3 and RPTOR; however such data need additional evidences. Not involved in ammonia-induced autophagy or in autophagic response of cerebellar granule neurons (CGN) to low potassium concentration. Plays a role early in neuronal differentiation and is required for granule cell axon formation: may govern axon formation via Ras-like GTPase signaling and through regulation of the Rab5-mediated endocytic pathways within developing axons.

Cellular Location

Cytoplasmic vesicle membrane; Peripheral membrane protein. Note=Localizes to pre-autophagosomal membrane

References

1.J Biol Chem. 2014 Aug 8;289(32):22306-18.2.Oncogene. 1999 Oct 21;18(43):5850-9.

Images

