

VEGFA Antibody

Purified Mouse Monoclonal Antibody Catalog # AO2228a

Product Information

Application WB, ICC, E **Primary Accession** P15692 Reactivity Human Host Mouse Monoclonal Clonality **Clone Names** 6G5B5 Isotype IgG1 **Calculated MW** 43597

Description This gene is a member of the PDGF/VEGF growth factor family. It encodes a

heparin-binding protein, which exists as a disulfide-linked homodimer. This growth factor induces proliferation and migration of vascular endothelial cells, and is essential for both physiological and pathological angiogenesis. Disruption of this gene in mice resulted in abnormal embryonic blood vessel formation. This gene is upregulated in many known tumors and its expression is correlated with tumor stage and progression. Elevated levels of this protein are found in patients with POEMS syndrome, also known as Crow-Fukase

syndrome. Allelic variants of this gene have been associated with

microvascular complications of diabetes 1 (MVCD1) and atherosclerosis. Alternatively spliced transcript variants encoding different isoforms have been described. There is also evidence for alternative translation initiation from upstream non-AUG (CUG) codons resulting in additional isoforms. A recent study showed that a C-terminally extended isoform is produced by use of an alternative in-frame translation termination codon via a stop codon

readthrough mechanism, and that this isoform is antiangiogenic. Expression of some isoforms derived from the AUG start codon is regulated by a small upstream open reading frame, which is located within an internal ribosome

entry site.

Immunogen Purified recombinant fragment of human VEGFA (AA: 207-371) expressed in E.

Coli.

Formulation Purified antibody in PBS with 0.05% sodium azide

Additional Information

Gene ID 7422

Other Names Vascular endothelial growth factor A, VEGF-A, Vascular permeability factor,

VPF, VEGFA, VEGF

Dilution WB~~1/500 - 1/2000 ICC~~N/A E~~1/10000

Storage Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions VEGFA Antibody is for research use only and not for use in diagnostic or

therapeutic procedures.

Protein Information

Name VEGFA

Synonyms VEGF

Function [N-VEGF]: Participates in the induction of key genes involved in the response

to hypoxia and in the induction of angiogenesis such as HIF1A

(PubMed:35455969). Involved in protecting cells from hypoxia- mediated cell

death (By similarity).

Cellular Location [N-VEGF]: Cytoplasm. Nucleus. Note=Cytoplasmic in normoxic conditions and

localizes to the nucleus under hypoxic conditions [Isoform L-VEGF189]: Endoplasmic reticulum. Golgi apparatus. Secreted, extracellular space,

extracellular matrix [Isoform VEGF165]: Secreted

Tissue Location Higher expression in pituitary tumors than the pituitary gland. [Isoform

VEGF165]: Widely expressed. [Isoform VEGF206]: Not widely expressed.

References

1.Oncol Rep. 2014 Dec;32(6):2359-64. 2.Eur J Cancer. 2014 Nov;50(16):2855-65.

Images

