10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ALDH2 Antibody

Purified Mouse Monoclonal Antibody
Catalog # AO2295a

Product Information

Application	WB, IHC, FC, ICC, E
Primary Accession	P05091
Reactivity	Human, Mouse, Rat
Host	Mouse
Clonality	Monoclonal
Clone Names	4G6A3
Isotype	IgG1
Calculated MW	56381
Description	This protein belongs to the aldehyde dehydrogenase family of proteins. Aldehyde dehydrogenase is the second enzyme of the major oxidative pathway of alcohol metabolism. Two major liver isoforms of aldehyde dehydrogenase, cytosolic and mitochondrial, can be distinguished by their electrophoretic mobilities, kinetic properties, and subcellular localizations. Most Caucasians have two major isozymes, while approximately 50% of Orientals have the cytosolic isozyme but not the mitochondrial isozyme. A remarkably higher frequency of acute alcohol intoxication among Orientals than among Caucasians could be related to the absence of a catalytically active form of the mitochondrial isozyme. The increased exposure to acetaldehyde in individuals with the catalytically inactive form may also confer greater susceptibility to many types of cancer. This gene encodes a mitochondrial isoform, which has a low Km for acetaldehydes, and is localized in mitochondrial matrix. Alternative splicing results in multiple transcript variants encoding distinct isoforms.
Immunogen	Purified recombinant fragment of human ALDH2 (AA: 317-517) expressed in E. Coli.
Formulation	Purified antibody in PBS with 0.05% sodium azide

Additional Information

Gene ID	217
Other Names	Aldehyde dehydrogenase, mitochondrial, 1.2.1.3, ALDH class 2, ALDH-E2, ALDHI, ALDH2, ALDM
Dilution	WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 FC~~1/200 - 1/400 ICC~~N/A E~~1/10000
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions	ALDH2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ALDH2 (HGNC:404)
Synonyms	ALDM
Function	Mitochondrial aldehyde dehydrogenase with broad substrate specificity which participates in the detoxification of ethanol-derived acetaldehyde and endogenous aldehydes, including those arising from lipid peroxidation (PubMed:16956664, PubMed:22339434, PubMed:33355142, PubMed:39075523). The reaction mechanism involves a nucleophilic attack on the aldehyde substrate, resulting in formation of a thiohemiacetal intermediate. Subsequent hydride transfer to NAD+, yields a thioester intermediate which is hydrolyzed to the carboxylic acid product through activation of a water molecule by the catalytic glutamate (PubMed:19506075). In vivo, ALDH2 is required for the clearance of formaldehyde, a cytotoxic and carcinogenic endogenous metabolite that induces DNA damage (PubMed:33355142). It also catalyzes the oxidation of 3,4-dihydroxyphenylacetaldehyde (DOPAL), a reactive intermediate of dopamine catabolism, to 3,4-dihydroxyphenylacetate (DOPAC) (PubMed:16956664). ALDH2 may also catalyze the oxidation of 4-aminobutanal (gamma-aminobutyraldehyde) to 4-aminobutanoate (GABA), representing an alternative route for GABA biosynthesis from polyamine catabolism (By similarity). In vitro, ALDH2 catalyzes the oxidation of acrolein, malondialdehyde, and 4-hydroxynonenal (4-HNE) (PubMed:22339434). It also exhibits esterase activity that is enhanced by NAD+ but does not require it, although the physiological relevance of this activity remains unclear (PubMed:19506075).
Cellular Location	Mitochondrion

References

1.Eur Heart J. 2012 Jul;33(13):1606-14. 2.Clin Toxicol (Phila). 2012 Apr;50(4):242-9.

Images





Figure 1: Western blot analysis using ALDH2 mAb against human ALDH2 recombinant protein. (Expected MW is 47.4 kDa)

Figure 2: Western blot analysis using ALDH2 mAb against HEK293 (1) and ALDH2 (AA: 317-517)-hIgGFc transfected HEK293 (2) cell lysate.






Figure 3: Western blot analysis using ALDH2 mouse mAb against HepG2 (1), A549 (2) cell lysate, and Rat live (3), Mouse liver (4), Mouse brain (5) tissue lysate.




Figure 4: Immunofluorescence analysis of HepG2 cells using ALDH2 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye.




Figure 5: Flow cytometric analysis of HeLa cells using ALDH2 mouse mAb (green) and negative control (red).




Figure 6: Immunohistochemical analysis of paraffin-embedded cervical cancer tissues using ALDH2 mouse mAb with DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.