

# Mouse Monoclonal Antibody to Rab6b

Purified Mouse Monoclonal Antibody

Catalog # AO2354a

## Product Information

---

<b>Application</b>	WB, IHC, E
<b>Primary Accession</b>	<a href="#">Q9NRW1</a>
<b>Reactivity</b>	Human, Rat
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	6D12G9
<b>Isotype</b>	Mouse IgG1
<b>Calculated MW</b>	23462
<b>Description</b>	RAB6B (RAB6B, Member RAS Oncogene Family) is a Protein Coding gene. Among its related pathways are Sertoli-Sertoli Cell Junction Dynamics. GO annotations related to this gene include GTP binding and GTPase activity. An important paralog of this gene is RAB41.;
<b>Immunogen</b>	Purified recombinant fragment of human Rab6b (AA: 95-208) expressed in E. Coli.
<b>Formulation</b>	Purified antibody in PBS with 0.05% sodium azide
<b>Application Note</b>	ELISA: 1/10000; WB: 1/500 - 1/2000; IHC: 1/200 - 1/1000;

## Additional Information

---

<b>Gene ID</b>	51560
<b>Other Names</b>	Ras-related protein Rab-6B, RAB6B
<b>Dilution</b>	WB~~1:1000 IHC~~1:100~500 E~~N/A
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	Mouse Monoclonal Antibody to Rab6b is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	RAB6B ( <a href="#">HGNC:14902</a> )
<b>Function</b>	The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with

membranes. Rabs cycle between active GTP-bound and inactive GDP-bound states. In their active state, drive transport of vesicular carriers from donor organelles to acceptor organelles to regulate the membrane traffic that maintains organelle identity and morphology (By similarity). Recruits VPS13B to the Golgi membrane (PubMed:[25492866](#)). Regulates the compacted morphology of the Golgi (PubMed:[26209634](#)). Seems to have a role in retrograde membrane traffic at the level of the Golgi complex. May function in retrograde transport in neuronal cells (PubMed:[17707369](#)). Plays a role in neuron projection development (PubMed:[25492866](#)).

## Cellular Location

Golgi apparatus membrane; Lipid-anchor. Endoplasmic reticulum-Golgi intermediate compartment Cytoplasmic vesicle. Note=Colocalizes with BICD1 at vesicular structures that align along microtubules


## Tissue Location

Predominantly expressed in brain.


## References

1.Exp Cell Res. 2007 Oct 1;313(16):3408-20. ; 2.J Cell Sci. 2000 Aug;113 ( Pt 15):2725-35.;

## Images




Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)





Western blot analysis using Rab6b mAb against human Rab6b (AA: 95-208) recombinant protein. (Expected MW is 38.7 kDa)

Western blot analysis using Rab6b mAb against HEK293 (1) and Rab6b (AA: 95-208)-hIgGfc transfected HEK293 (2) cell lysate.



Western blot analysis using Rab6b mouse mAb against C6 (1), SK-N-SH (2), HT-29 (3), PC-12 (4), and C6 (5) cell lysate.



Immunohistochemical analysis of paraffin-embedded bladder cancer tissues using Rab6b mouse mAb with DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.