

Mouse Monoclonal Antibody to MITF

Purified Mouse Monoclonal Antibody

Catalog # AO2420a

Product Information

Application	WB, FC, E
Primary Accession	O75030
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Clone Names	8F1G5
Isotype	Mouse IgG1
Calculated MW	58795
Description	This gene encodes a transcription factor that contains both basic helix-loop-helix and leucine zipper structural features. It regulates the differentiation and development of melanocytes retinal pigment epithelium and is also responsible for pigment cell-specific transcription of the melanogenesis enzyme genes. Heterozygous mutations in the this gene cause auditory-pigmentary syndromes, such as Waardenburg syndrome type 2 and Tietz syndrome. Alternatively spliced transcript variants encoding different isoforms have been identified.;
Immunogen	Purified recombinant fragment of human MITF (AA: 1-114) expressed in E. Coli.
Formulation	Purified antibody in PBS with 0.05% sodium azide
Application Note	ELISA: 1/10000; WB: 1/500 - 1/2000; FCM: 1/200 - 1/400

Additional Information

Gene ID	4286
Other Names	MI; WS2; CMM8; WS2A; bHLHe32
Dilution	WB~~1:1000 FC~~1:10~50 E~~N/A
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Mouse Monoclonal Antibody to MITF is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Name	MITF {ECO:0000303 PubMed:8069297, ECO:0000312 HGNC:HGNC:7105}
Function	Transcription factor that acts as a master regulator of melanocyte survival and differentiation as well as melanosome biogenesis (PubMed: 10587587 , PubMed: 22647378 , PubMed: 27889061 , PubMed: 9647758). Binds to M-boxes (5'-TCATGTG-3') and symmetrical DNA sequences (E-boxes) (5'-CACGTG-3') found in the promoter of pigmentation genes, such as tyrosinase (TYR) (PubMed: 10587587 , PubMed: 22647378 , PubMed: 27889061 , PubMed: 9647758). Involved in the cellular response to amino acid availability by acting downstream of MTOR: in the presence of nutrients, MITF phosphorylation by MTOR promotes its inactivation (PubMed: 36608670). Upon starvation or lysosomal stress, inhibition of MTOR induces MITF dephosphorylation, resulting in transcription factor activity (PubMed: 36608670). Plays an important role in melanocyte development by regulating the expression of tyrosinase (TYR) and tyrosinase-related protein 1 (TYRP1) (PubMed: 10587587 , PubMed: 22647378 , PubMed: 27889061 , PubMed: 9647758). Plays a critical role in the differentiation of various cell types, such as neural crest-derived melanocytes, mast cells, osteoclasts and optic cup-derived retinal pigment epithelium (PubMed: 10587587 , PubMed: 22647378 , PubMed: 27889061 , PubMed: 9647758).
Cellular Location	Nucleus. Cytoplasm. Lysosome membrane Note=When nutrients are present, recruited to the lysosomal membrane via association with GDP-bound RagC/RRAGC (or RagD/RRAGD): it is then phosphorylated by MTOR (PubMed:23401004, PubMed:36608670) Phosphorylation by MTOR promotes ubiquitination and degradation (PubMed:36608670). Conversely, inhibition of mTORC1, starvation and lysosomal disruption, promotes dephosphorylation and translocation to the nucleus (PubMed:36608670). Phosphorylation by MARK3/cTAK1 promotes association with 14-3-3/YWHA adapters and retention in the cytosol (PubMed:16822840).
Tissue Location	Expressed in melanocytes (at protein level). [Isoform C2]: Expressed in the kidney and retinal pigment epithelium. [Isoform H2]: Expressed in the kidney. [Isoform Mdel]: Expressed in melanocytes.

References


1.Proc Natl Acad Sci U S A. 2015 Feb 3;112(5):E420-9. ; 2.Cell Mol Life Sci. 2015 Apr;72(7):1249-60.;

Images




Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)



Western blot analysis using MITF mAb against human MITF (AA: 1-114) recombinant protein. (Expected MW is 38.9 kDa)



Western blot analysis using MITF mAb against HEK293 (1) and MITF (AA: 1-114)-hIgGfc transfected HEK293 (2) cell lysate.



Western blot analysis using MITF mouse mAb against PANC-1 (1) and A549 (2) cell lysate.



Flow cytometric analysis of HeLa cells using MITF mouse mAb (green) and negative control (red).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.