10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Mouse Monoclonal Antibody to IDH1

Purified Mouse Monoclonal Antibody
Catalog # AO2436a

## Product Information

| | |
|---|---|
| **Application** | WB, FC, ICC, E |
| **Primary Accession** | O75874 |
| **Reactivity** | Human, Mouse, Monkey |
| **Host** | Mouse |
| **Clonality** | Monoclonal |
| **Clone Names** | 7G8A1 |
| **Isotype** | Mouse IgG1 |
| **Calculated MW** | 46659 |
| **Description** | Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. Each NADP(+)-dependent isozyme is a homodimer. The protein encoded by this gene is the NADP(+)-dependent isocitrate dehydrogenase found in the cytoplasm and peroxisomes. It contains the PTS-1 peroxisomal targeting signal sequence. The presence of this enzyme in peroxisomes suggests roles in the regeneration of NADPH for intraperoxisomal reductions, such as the conversion of 2, 4-dienoyl-CoAs to 3-enoyl-CoAs, as well as in peroxisomal reactions that consume 2-oxoglutarate, namely the alpha-hydroxylation of phytanic acid. The cytoplasmic enzyme serves a significant role in cytoplasmic NADPH production. Alternatively spliced transcript variants encoding the same protein have been found for this gene.; |
| **Immunogen** | Purified recombinant fragment of human IDH1 (AA: 156-298) expressed in E. Coli. |
| **Formulation** | Purified antibody in PBS with 0.05% sodium azide |
| **Application Note** | ELISA: 1/10000; WB: 1/500 - 1/2000; ICC: 1/50 - 1/250; FCM: 1/200 - 1/400 |

## Additional Information

| | |
|---|---|
| **Gene ID** | 3417 |
| **Other Names** | IDH; IDP; IDCD; IDPC; PICD; HEL-216; HEL-S-26 |
| **Dilution** | WB~~1:1000 FC~~1:10~50 ICC~~N/A E~~N/A |
| **Storage** | Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store |

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** Mouse Monoclonal Antibody to IDH1 is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

**Name** IDH1

**Synonyms** PICD


**Function** Catalyzes the NADP(+)-dependent oxidative decarboxylation of isocitrate (D-threo-isocitrate) to 2-ketoglutarate (2-oxoglutarate), which is required by other enzymes such as the phytanoyl-CoA dioxygenase (PubMed:10521434, PubMed:19935646). Plays a critical role in the generation of NADPH, an important cofactor in many biosynthesis pathways (PubMed:10521434). May act as a corneal epithelial crystallin and may be involved in maintaining corneal epithelial transparency (By similarity).

**Cellular Location** Cytoplasm, cytosol. Peroxisome


## References

1.Cancer Cell. 2015 Dec 14;28(6):773-84. ; 2.Int J Cancer. 2015 Sep 1;137(5):1058-65. ;

## Images






Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)





Western blot analysis using IDH1 mAb against human IDH1 (AA: 156-298) recombinant protein. (Expected MW is 41.8 kDa)

Western blot analysis using IDH1 mAb against HEK293 (1) and IDH1 (AA: 156-298)-hIgGFc transfected HEK293 (2) cell lysate.





Western blot analysis using IDH1 mouse mAb against HepG2 (1), NIH/3T3 (2), C2C12 (3), COS7 (4), and SW480 (5) cell lysate.



Immunofluorescence analysis of Hela cells using IDH1 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin. Secondary antibody from Fisher




Flow cytometric analysis of Hela cells using IDH1 mouse mAb (green) and negative control (red).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.