

# Mouse Monoclonal Antibody to KRT19

Purified Mouse Monoclonal Antibody

Catalog # AO2460a

## Product Information

---

<b>Application</b>	WB, IHC, E
<b>Primary Accession</b>	<a href="#">P08727</a>
<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	8H1A8
<b>Isotype</b>	Mouse IgG2a
<b>Calculated MW</b>	44106
<b>Description</b>	The protein encoded by this gene is a member of the keratin family. The keratins are intermediate filament proteins responsible for the structural integrity of epithelial cells and are subdivided into cytokeratins and hair keratins. The type I cytokeratins consist of acidic proteins which are arranged in pairs of heterotypic keratin chains. Unlike its related family members, this smallest known acidic cytokeratin is not paired with a basic cytokeratin in epithelial cells. It is specifically expressed in the periderm, the transiently superficial layer that envelopes the developing epidermis. The type I cytokeratins are clustered in a region of chromosome 17q12-q21.;
<b>Immunogen</b>	Purified recombinant fragment of human KRT19 (AA: 274-400) expressed in E. Coli.
<b>Formulation</b>	Purified antibody in PBS with 0.05% sodium azide
<b>Application Note</b>	ELISA: 1/10000; WB: 1/500 - 1/2000; IHC: 1/200 - 1/1000;

## Additional Information

---

<b>Gene ID</b>	3880
<b>Other Names</b>	K19; CK19; K1CS
<b>Dilution</b>	WB~~1:1000 IHC~~1:100~500 E~~N/A
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	Mouse Monoclonal Antibody to KRT19 is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information


---

<b>Name</b>	KRT19
<b>Function</b>	Involved in the organization of myofibers. Together with KRT8, helps to link the contractile apparatus to dystrophin at the costameres of striated muscle.
<b>Tissue Location</b>	Expressed in a defined zone of basal keratinocytes in the deep outer root sheath of hair follicles. Also observed in sweat gland and mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, ectocervical epithelium (at protein level). Expressed in epidermal basal cells, in nipple epidermis and a defined region of the hair follicle. Also seen in a subset of vascular wall cells in both the veins and artery of human umbilical cord, and in umbilical cord vascular smooth muscle. Observed in muscle fibers accumulating in the costameres of myoplasm at the sarcolemma in structures that contain dystrophin and spectrin.


## References

1.Clin Cancer Res. 2015 Jul 1;21(13):3081-91. ; 2.Gut. 2014 Apr;63(4):674-85. ;

## Images




Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)





Western blot analysis using KRT19 mAb against human KRT19 (AA: 274-400) recombinant protein. (Expected MW is 40 kDa)


Western blot analysis using KRT19 mAb against HEK293 (1) and KRT19 (AA: 274-400)-hIgGFc transfected HEK293 (2) cell lysate.



Western blot analysis using KRT19 mouse mAb against MCF-7 (1), SW480 (2), SW620 (3), and HCT116 (4) cell lysate.



Immunohistochemical analysis of paraffin-embedded stomach cancer tissues using KRT19 mouse mAb with DAB staining.



Immunohistochemical analysis of paraffin-embedded endometrial cancer tissues using KRT19 mouse mAb with DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.