

Mouse Monoclonal Antibody to ANAPC10

Purified Mouse Monoclonal Antibody

Catalog # AO2468a

Product Information

| | |
|--------------------------|--|
| Application | WB, IHC, ICC, E |
| Primary Accession | Q9UM13 |
| Reactivity | Human |
| Host | Mouse |
| Clonality | Monoclonal |
| Clone Names | 3E9A9 |
| Isotype | Mouse IgG1 |
| Calculated MW | 21252 |
| Description | ANAPC10 is a core subunit of the anaphase-promoting complex (APC), or cyclosome, a ubiquitin protein ligase that is essential for progression through the cell cycle. APC initiates sister chromatid separation by ubiquitinating the anaphase inhibitor securin (PTTG1; MIM 604147) and triggers exit from mitosis by ubiquitinating cyclin B (CCNB1; MIM 123836), the activating subunit of cyclin-dependent kinase-1 (CDK1; MIM 116940) (summary by Wendt et al., 2001 [PubMed 11524682]).; |
| Immunogen | Purified recombinant fragment of human ANAPC10 (AA: 1-185) expressed in E. Coli. |
| Formulation | Purified antibody in PBS with 0.05% sodium azide |
| Application Note | ELISA: 1/10000; WB: 1/500 - 1/2000; IHC: 1/200 - 1/1000; ICC: 1/200 - 1/1000; |

Additional Information

| | |
|--------------------|--|
| Gene ID | 10393 |
| Other Names | DOC1; APC10 |
| Dilution | WB~~1:1000 IHC~~1:100~500 ICC~~N/A E~~N/A |
| Storage | Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| Precautions | Mouse Monoclonal Antibody to ANAPC10 is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information

| | |
|-------------|---------|
| Name | ANAPC10 |
|-------------|---------|

Synonyms

APC10


Function

Component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle (PubMed:[18485873](#)). The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains (PubMed:[18485873](#)). The APC/C complex catalyzes assembly of branched 'Lys-11'-/'Lys-48'-linked branched ubiquitin chains on target proteins (PubMed:[29033132](#)).


References

1.BMC Cell Biol. 2004 May 16;5:20. ; 2.Nat Struct Biol. 2001 Sep;8(9):784-8.;


Images




Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)




Western blot analysis using ANAPC10 mAb against human ANAPC10 (AA: 1-185) recombinant protein. (Expected MW is 47.2 kDa)




Western blot analysis using ANAPC10 mAb against HEK293 (1) and ANAPC10 (AA: 1-185)-hIgGfc transfected HEK293 (2) cell lysate.




Western blot analysis using ANAPC10 mouse mAb against Jurkat (1), MCF-7 (2), SK-Br-3 (3), A431 (4), HEK293 (5), A549 (6), and SPC-A-1 (7) cell lysate.



Immunofluorescence analysis of HeLa cells using ANAPC10 mouse mAb. Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.



Immunofluorescence analysis of HeLa cells using ANAPC10 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin. Secondary antibody from Fisher



Immunohistochemical analysis of paraffin-embedded esophageal cancer tissues using ANAPC10 mouse mAb with DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.