

Mouse Monoclonal Antibody to ADAMTS1

Purified Mouse Monoclonal Antibody Catalog # AO2479a

Product Information

Application WB, E **Primary Accession** Q9UHI8 Reactivity Human Host Mouse Monoclonal Clonality **Clone Names** 1G5D5 Isotype Mouse IgG1 **Calculated MW** 105358

Description This gene encodes a member of the ADAMTS (a disintegrin and

metalloproteinase with thrombospondin motif) protein family. Members of the family share several distinct protein modules, including a propeptide region, a metalloproteinase domain, a disintegrin-like domain, and a thrombospondin type 1 (TS) motif. Individual members of this family differ in the number of C-terminal TS motifs, and some have unique C-terminal domains. The protein encoded by this gene contains two disintegrin loops and three C-terminal TS motifs and has anti-angiogenic activity. The expression of this gene may be associated with various inflammatory processes as well as development of cancer cachexia. This gene is likely to be necessary for normal growth, fertility, and organ morphology and function.;

Immunogen Purified recombinant fragment of human ADAMTS1 (AA: 858-960) expressed

in E. Coli.

Formulation Purified antibody in PBS with 0.05% sodium azide

Application Note ELISA: 1/10000; WB: 1/500 - 1/2000;

Additional Information

Gene ID 9510

Other Names C3-C5; METH1

Dilution WB~~1:1000 E~~N/A

Storage Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

PrecautionsMouse Monoclonal Antibody to ADAMTS1 is for research use only and not for

use in diagnostic or therapeutic procedures.

Protein Information

Name ADAMTS1

Synonyms KIAA1346, METH1

Function Metalloprotease which cleaves aggrecan, a cartilage proteoglycan, at the

'1938-Glu-|-Leu-1939' site (within the chondroitin sulfate attachment domain), and may be involved in its turnover (By similarity). Also cleaves

COMP (PubMed: <u>39672391</u>). Has angiogenic inhibitor activity

(PubMed: 10438512). May play a critical role in follicular rupture (By


similarity).

Cellular Location Secreted, extracellular space, extracellular matrix


References

1.Mol Cancer. 2013 Jan 5;12:2.; 2.Cancer Sci. 2012 Oct;103(10):1889-97.;

Images





Black line: Control Antigen (100 ng); Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)



Western blot analysis using ADAMTS1 mAb against human ADAMTS1 (AA: 858-960) recombinant protein. (Expected MW is 38.3 kDa)

Western blot analysis using ADAMTS1 mAb against HEK293 (1) and ADAMTS1 (AA: 858-960)-hIgGFc transfected HEK293 (2) cell lysate.





Western blot analysis using ADAMTS1 mouse mAb against Hela (1) and SK-Br-3 (2) cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.