

NAA10

Purified Mouse Monoclonal Antibody

Catalog # AO2505a

Product Information

Application	WB, IHC, ICC, E
Primary Accession	P41227
Reactivity	Human, Mouse, Monkey
Host	Mouse
Clonality	Monoclonal
Clone Names	3G3E9
Isotype	Mouse IgG1
Calculated MW	26459
Immunogen	Purified recombinant fragment of human NAA10 (AA: 111-235) expressed in E. Coli.
Formulation	Purified antibody in PBS with 0.05% sodium azide

Additional Information

Gene ID	8260
Other Names	TE2; ARD1; NATD; ARD1A; ARD1P; OGDNS; DXS707; MCOPS1
Dilution	WB~~ 1/500 - 1/2000 IHC~~1:100~500 ICC~~ 1/200 - 1/1000 E~~ 1/10000
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	NAA10 is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	NAA10
Synonyms	ARD1, ARD1A, TE2
Function	Catalytic subunit of N-terminal acetyltransferase complexes which display alpha (N-terminal) acetyltransferase activity (PubMed: 15496142 , PubMed: 19420222 , PubMed: 19826488 , PubMed: 20145209 , PubMed: 20154145 , PubMed: 25489052 , PubMed: 27708256 , PubMed: 29754825 , PubMed: 32042062). Acetylates amino termini that are devoid of initiator methionine (PubMed: 19420222). The alpha (N-terminal) acetyltransferase activity may be important for vascular, hematopoietic and

neuronal growth and development. Without NAA15, displays epsilon (internal) acetyltransferase activity towards HIF1A, thereby promoting its degradation (PubMed:[12464182](#)). Represses MYLK kinase activity by acetylation, and thus represses tumor cell migration (PubMed:[19826488](#)). Acetylates, and stabilizes TSC2, thereby repressing mTOR activity and suppressing cancer development (PubMed:[20145209](#)). Acetylates HSPA1A and HSPA1B at 'Lys-77' which enhances its chaperone activity and leads to preferential binding to co-chaperone HOPX (PubMed:[27708256](#)). Acetylates HIST1H4A (PubMed:[29754825](#)). Acts as a negative regulator of sister chromatid cohesion during mitosis (PubMed:[27422821](#)).

Cellular Location

Cytoplasm. Nucleus. Note=Also present in the free cytosolic and cytoskeleton-bound polysomes.

Tissue Location

Ubiquitous..

References

1.Gene. 2015 Aug 10;567(2):103-31.2.PLoS One. 2014 Aug 18;9(8):e105185.

Images




Figure 1:Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)




Figure 2:Western blot analysis using NAA10 mAb against human NAA10 (AA: 111-235) recombinant protein. (Expected MW is 47.2 kDa)

Figure 3:Western blot analysis using NAA10 mAb against HEK293 (1) and NAA10 (AA: 111-235)-hIgGfc transfected HEK293 (2) cell lysate.




Figure 4: Western blot analysis using NAA10 mouse mAb against HCT116 (1), COS7 (2), HEK293 (3), HL-60 (4), MCF-7 (5), Hela (6), NIH/3T3 (7), and C2C12 (8) cell lysate.




Figure 5: Immunofluorescence analysis of Hela cells using NAA10 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin. Secondary antibody from Fisher (Cat#: 35503)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.