

ALDH1A1

Purified Mouse Monoclonal Antibody

Catalog # AO2560a

Product Information

Application	WB, IHC, ICC, E
Primary Accession	P00352
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Clone Names	2G1G4
Isotype	Mouse IgG1
Calculated MW	54862
Immunogen	Purified recombinant fragment of human ALDH1A1 (AA: 1-110) expressed in E. Coli.
Formulation	Purified antibody in PBS with 0.05% sodium azide

Additional Information

Gene ID	216
Other Names	ALDC; ALDH1; HEL-9; HEL12; PUMB1; ALDH11; RALDH1; ALDH-E1; HEL-S-53e
Dilution	WB~~ 1/500 - 1/2000 IHC~~ 1/200 - 1/1000 ICC~~N/A E~~ 1/10000
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ALDH1A1 is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ALDH1A1 (HGNC:402)
Function	Cytosolic dehydrogenase that catalyzes the irreversible oxidation of a wide range of aldehydes to their corresponding carboxylic acid (PubMed: 12941160 , PubMed: 15623782 , PubMed: 17175089 , PubMed: 19296407 , PubMed: 25450233 , PubMed: 26373694). Functions downstream of retinol dehydrogenases and catalyzes the oxidation of retinaldehyde into retinoic acid, the second step in the oxidation of retinol/vitamin A into retinoic acid (By similarity). This pathway is crucial to control the levels of retinol and retinoic acid, two important molecules which excess can be teratogenic and cytotoxic (By similarity). Also oxidizes aldehydes resulting from lipid

peroxidation like (E)-4-hydroxynon-2-enal/HNE, malonaldehyde and hexanal that form protein adducts and are highly cytotoxic. By participating for instance to the clearance of (E)-4-hydroxynon-2-enal/HNE in the lens epithelium prevents the formation of HNE-protein adducts and lens opacification (PubMed:[12941160](#), PubMed:[15623782](#), PubMed:[19296407](#)). Also functions downstream of fructosamine-3-kinase in the fructosamine degradation pathway by catalyzing the oxidation of 3-deoxyglucosone, the carbohydrate product of fructosamine 3-phosphate decomposition, which is itself a potent glycating agent that may react with lysine and arginine side-chains of proteins (PubMed:[17175089](#)). Also has an aminobutyraldehyde dehydrogenase activity and is probably part of an alternative pathway for the biosynthesis of GABA/4-aminobutanoate in midbrain, thereby playing a role in GABAergic synaptic transmission (By similarity).

Cellular Location Cytoplasm, cytosol. Cell projection, axon {ECO:0000250 | UniProtKB:P24549}

Tissue Location Expressed by erythrocytes (at protein level).

References

1.Oncotarget. 2015 Dec 1;6(38):41360-9.2.Biomark Med. 2015;9(8):777-90.

Images




Figure 1:Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)




Figure 2:Western blot analysis using ALDH1A1 mAb against human ALDH1A1 (AA: 1-110) recombinant protein. (Expected MW is 38.4 kDa)

Figure 3:Western blot analysis using ALDH1A1 mAb against HEK293 (1) and ALDH1A1 (AA: 1-110)-hIgGfc transfected HEK293 (2) cell lysate.




Figure 4: Western blot analysis using ALDH1A1 mouse mAb against HepG2 (1) and A549 (2) cell lysate.






Figure 5: Immunohistochemical analysis of paraffin-embedded cervical cancer tissues using ALDH1A1 mouse mAb with DAB staining.



1/200 - 1/1000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.