

# HAVCR1

Purified Mouse Monoclonal Antibody

Catalog # AO2576a

## Product Information

---

<b>Application</b>	WB, IHC, ICC, E
<b>Primary Accession</b>	<a href="#">Q96D42</a>
<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	3A12E10
<b>Isotype</b>	Mouse IgG1
<b>Calculated MW</b>	39250
<b>Immunogen</b>	Purified recombinant fragment of human HAVCR1 (AA: 70-290) expressed in E. Coli.
<b>Formulation</b>	Purified antibody in PBS with 0.05% sodium azide

## Additional Information

---

<b>Gene ID</b>	26762
<b>Other Names</b>	TIM; KIM1; TIM1; CD365; HAVCR; KIM-1; TIM-1; TIMD1; TIMD-1; HAVCR-1
<b>Dilution</b>	WB~~ 1/500 - 1/2000 IHC~~1:100~500 ICC~~N/A E~~ 1/10000
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	HAVCR1 is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	HAVCR1
<b>Synonyms</b>	KIM1, TIM1, TIMD1
<b>Function</b>	Phosphatidylserine receptor that plays an important functional role in regulatory B-cells homeostasis including generation, expansion and suppressor functions (By similarity). As P- selectin/SELPLG ligand, plays a specialized role in activated but not naive T-cell trafficking during inflammatory responses (PubMed: <a href="#">24703780</a> ). Controls thereby T-cell accumulation in the inflamed central nervous system (CNS) and the induction of autoimmune disease (PubMed: <a href="#">24703780</a> ). Also regulates expression of

various anti- inflammatory cytokines and co-inhibitory ligands including IL10 (By similarity). Acts as a regulator of T-cell proliferation (By similarity). May play a role in kidney injury and repair (PubMed:[17471468](#)).

## Cellular Location

Cell membrane; Single-pass type I membrane protein

## Tissue Location

Widely expressed, with highest levels in kidney and testis. Expressed by activated CD4+ T-cells during the development of helper T-cells responses.

## References

1.Biomed Res Int. 2015;2015:854070. 2.Pediatr Res. 2015 Oct;78(4):430-5.

## Images




Figure 1:Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)




Figure 2:Western blot analysis using HAVCR1 mAb against human HAVCR1 (AA: 70-290) recombinant protein. (Expected MW is 37 kDa)





Figure 3:Western blot analysis using HAVCR1 mAb against HEK293 (1) and HAVCR1 (AA: 70-290)-hIgGfc transfected HEK293 (2) cell lysate.

Figure 4:Flow cytometric analysis of Hela cells using HAVCR1 mouse mAb (green) and negative control (red).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.