

# ATPIF1

Purified Mouse Monoclonal Antibody  
Catalog # AO2601a

## Product Information

---

<b>Application</b>	WB, IHC, ICC, E
<b>Primary Accession</b>	<a href="#">Q9UII2</a>
<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	1F3B8
<b>Isotype</b>	Mouse IgG2b
<b>Calculated MW</b>	12249
<b>Immunogen</b>	Purified recombinant fragment of human ATPIF1 (AA: 1-106) expressed in E. Coli.
<b>Formulation</b>	Purified antibody in PBS with 0.05% sodium azide

## Additional Information

---

<b>Gene ID</b>	93974
<b>Other Names</b>	IP; ATP1; ATPIP
<b>Dilution</b>	WB~~ 1/500 - 1/2000 IHC~~ 1/200 - 1/1000 ICC~~ 1/100 - 1/500 E~~ 1/10000
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	ATPIF1 is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	ATP5IF1 ( <a href="#">HGNC:871</a> )
<b>Synonyms</b>	ATPI, ATPIF1
<b>Function</b>	Endogenous F(1)F(o)-ATPase inhibitor limiting ATP depletion when the mitochondrial membrane potential falls below a threshold and the F(1)F(o)-ATP synthase starts hydrolyzing ATP to pump protons out of the mitochondrial matrix. Required to avoid the consumption of cellular ATP when the F(1)F(o)-ATP synthase enzyme acts as an ATP hydrolase. Indirectly acts as a regulator of heme synthesis in erythroid tissues: regulates heme synthesis by modulating the mitochondrial pH and redox potential, allowing

FECH to efficiently catalyze the incorporation of iron into protoporphyrin IX to produce heme.

## Cellular Location

Mitochondrion.

## References

1.Pathobiology. 2015;82(5):224-32.2.Cell Rep. 2014 Apr 10;7(1):27-34.

## Images




Figure 1: Black line: Control Antigen (100 ng); Purple line: Antigen (10 ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)




Figure 2: Western blot analysis using ATPIF1 mAb against human ATPIF1 (AA: 1-106) recombinant protein. (Expected MW is 38.2 kDa)




Figure 3: Western blot analysis using ATPIF1 mAb against HEK293 (1) and ATPIF1 (AA: 1-106)-hIgGfc transfected HEK293 (2) cell lysate.

Figure 5: Flow cytometric analysis of Hela cells using ATPIF1 mouse mAb (green) and negative control (red).





Figure 4: Immunofluorescence analysis of HeLa cells using ATPIF1 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor- 555 phalloidin. Secondary antibody from Fisher (Cat#: 35503)




Figure 6: Immunohistochemical analysis of paraffin-embedded lung cancer tissues using ATPIF1 mouse mAb with DAB staining.




Figure 7: Immunohistochemical analysis of paraffin-embedded cervical cancer tissues using ATPIF1 mouse mAb with DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.