10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

TNFSF11

Purified Mouse Monoclonal Antibody
Catalog # AO2634a

Product Information

Application	WB, IHC, ICC, E
Primary Accession	O14788
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Clone Names	3H8A7
Isotype	Mouse IgG1
Calculated MW	35478
Immunogen	Purified recombinant fragment of human TNFSF11 (AA: 74-308) expressed in E. Coli.
Formulation	Purified antibody in PBS with 0.05% sodium azide

Additional Information

Gene ID	8600
Other Names	CD254; ODF; OPGL; sOdf; OPTB2; RANKL; TNLG6B; TRANCE; hRANKL2
Dilution	WB~~ 1/500 - 1/2000 IHC~~ 1/200 - 1/1000 ICC~~N/A E~~ 1/10000
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TNFSF11 is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TNFSF11 (HGNC:11926)
Synonyms	OPGL, RANKL, TRANCE
Function	Cytokine that binds to TNFRSF11B/OPG and to TNFRSF11A/RANK. Osteoclast differentiation and activation factor. Augments the ability of dendritic cells to stimulate naive T-cell proliferation. May be an important regulator of interactions between T-cells and dendritic cells and may play a role in the regulation of the T-cell-dependent immune response. May also play an important role in enhanced bone-resorption in humoral hypercalcemia of malignancy (PubMed:22664871). Induces osteoclastogenesis by activating

multiple signaling pathways in osteoclast precursor cells, chief among which is induction of long lasting oscillations in the intracellular concentration of Ca (2+) resulting in the activation of NFATC1, which translocates to the nucleus and induces osteoclast-specific gene transcription to allow differentiation of osteoclasts. During osteoclast differentiation, in a TMEM64 and ATP2A2-dependent manner induces activation of CREB1 and mitochondrial ROS generation necessary for proper osteoclast generation (By similarity).

Cellular Location

[Isoform 1]: Cell membrane; Single-pass type II membrane protein [Isoform 2]: Cytoplasm.

Tissue Location

Highest in the peripheral lymph nodes, weak in spleen, peripheral blood Leukocytes, bone marrow, heart, placenta, skeletal muscle, stomach and thyroid

References

1.Breast Cancer Res. 2015 Feb 21;17:24.2.Immunobiology. 2015 May;220(5):692-700.

Images




Figure 1:Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)





Figure 2:Western blot analysis using TNFSF11 mAb against human TNFSF11 (AA: 74-308) recombinant protein. (Expected MW is 52.6 kDa)

Figure 3:Western blot analysis using TNFSF11 mAb against HEK293 (1) and TNFSF11 (AA: 74-308)-hIgGFc transfected HEK293 (2) cell lysate.






Figure 4:Western blot analysis using TNFSF11 mouse mAb against U937 (1), HL-60 (2), Raji (3), and Ramos (4) cell lysate.




Figure 5:Flow cytometric analysis of Hela cells using TNFSF11 mouse mAb (green) and negative control (red).




Figure 6:Immunohistochemical analysis of paraffin-embedded bladder cancer tissues using TNFSF11 mouse mAb with DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.