

# CD152

Purified Mouse Monoclonal Antibody  
Catalog # AO2681a

## Product Information

---

<b>Application</b>	WB, IHC, ICC, E
<b>Primary Accession</b>	<a href="#">P16410</a>
<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	8B3F8
<b>Isotype</b>	Mouse IgG1
<b>Calculated MW</b>	24656
<b>Immunogen</b>	Purified recombinant fragment of human CD152 (AA: extra 36-161) expressed in E. Coli.
<b>Formulation</b>	Purified antibody in PBS with 0.05% sodium azide

## Additional Information

---

<b>Gene ID</b>	1493
<b>Other Names</b>	CTLA4; CD; GSE; GRD4; ALPS5; CTLA-4; IDDM12; CELIAC3
<b>Dilution</b>	WB~~ 1/500 - 1/2000 IHC~~1:100~500 ICC~~N/A E~~ 1/10000
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	CD152 is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	CTLA4
<b>Synonyms</b>	CD152
<b>Function</b>	Inhibitory receptor acting as a major negative regulator of T-cell responses. The affinity of CTLA4 for its natural B7 family ligands, CD80 and CD86, is considerably stronger than the affinity of their cognate stimulatory coreceptor CD28.
<b>Cellular Location</b>	Cell membrane; Single-pass type I membrane protein. Note=Exists primarily an intracellular antigen whose surface expression is tightly regulated by

restricted trafficking to the cell surface and rapid internalization

## Tissue Location

Widely expressed with highest levels in lymphoid tissues. Detected in activated T-cells where expression levels are 30- to 50-fold less than CD28, the stimulatory coreceptor, on the cell surface following activation.

## References

1.Asian Pac J Cancer Prev. 2016;17(8):3785-91.2.Eur J Cancer. 2015 Nov;51(17):2689-97.

## Images




Figure 4:Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)




Figure 2:Western blot analysis using CD152 mAb against human CD152 (AA: extra 36-161) recombinant protein. (Expected MW is 39.5 kDa)





Figure 3:Western blot analysis using CD152 mAb against HEK293 (1) and CD152 (AA: extra 36-161)-hIgGfc transfected HEK293 (2) cell lysate.

Figure 4:Flow cytometric analysis of Ramos cells using CD152 mouse mAb (green) and negative control (red).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.