

# PLXNA1

Purified Mouse Monoclonal Antibody  
Catalog # AO2684a

## Product Information

---

<b>Application</b>	WB, IHC, ICC, E
<b>Primary Accession</b>	<a href="#">Q9UIW2</a>
<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	7B8H9
<b>Isotype</b>	Mouse IgG2b
<b>Calculated MW</b>	211067
<b>Immunogen</b>	Purified recombinant fragment of human PLXNA1 (AA: 1100-1200) expressed in E. Coli.
<b>Formulation</b>	Purified antibody in PBS with 0.05% sodium azide

## Additional Information

---

<b>Gene ID</b>	5361
<b>Other Names</b>	NOV; NOVP; PLXN1; PLEXIN-A1
<b>Dilution</b>	WB~~ 1/500 - 1/2000 IHC~~1:100~500 ICC~~N/A E~~ 1/10000
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	PLXNA1 is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	PLXNA1 ( <a href="#">HGNC:9099</a> )
<b>Synonyms</b>	NOV, PLXN1
<b>Function</b>	Coreceptor for SEMA3A, SEMA3C, SEMA3F and SEMA6D. Necessary for signaling by class 3 semaphorins and subsequent remodeling of the cytoskeleton. Plays a role in axon guidance, invasive growth and cell migration. Class 3 semaphorins bind to a complex composed of a neuropilin and a plexin. The plexin modulates the affinity of the complex for specific semaphorins, and its cytoplasmic domain is required for the activation of down-stream signaling events in the cytoplasm. Acts as coreceptor of TREM2

for SEMA6D in dendritic cells and is involved in the generation of immune responses and skeletal homeostasis.

Cellular Location	Cell membrane {ECO:0000250   UniProtKB:P70206}; Single-pass type I membrane protein
Tissue Location	Detected in fetal brain, lung, liver and kidney.

References

1.PLoS One. 2016 Mar 10;11(3):e0149833. 2.World J Gastroenterol. 2007 Dec 28;13(48):6558-61.

Images




Figure 1:Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)




Figure 2:Western blot analysis using PLXNA1 mAb against human PLXNA1 (AA: 1100-1200) recombinant protein. (Expected MW is 36.8 kDa)





Figure 3:Western blot analysis using PLXNA1 mAb against HEK293 (1) and PLXNA1 (AA: 1100-1200)-hIgGfc transfected HEK293 (2) cell lysate.

Figure 4:Flow cytometric analysis of Hela cells using PLXNA1 mouse mAb (green) and negative control (red).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.