

# PD1

Purified Mouse Monoclonal Antibody  
Catalog # AO2750a

## Product Information

---

|                          |  |
|--------------------------|--|
| <b>Application</b>       | WB, IHC, ICC, E  |
| <b>Primary Accession</b> | <a href="#">Q15116</a>   |
| <b>Reactivity</b>        | Human  |
| <b>Host</b>              | Mouse  |
| <b>Clonality</b>         | Monoclonal   |
| <b>Clone Names</b>       | 1D6F6  |
| <b>Isotype</b>           | Mouse IgG1   |
| <b>Calculated MW</b>     | 31647  |
| <b>Immunogen</b>         | Purified recombinant fragment of human PD1 (AA: extra 21-170) expressed in HEK293 cells. |
| <b>Formulation</b>       | Purified antibody in PBS with 0.05% sodium azide   |

## Additional Information

---

|                    |  |
|--------------------|--|
| <b>Gene ID</b>     | 5133   |
| <b>Other Names</b> | PDCD1; PD-1; CD279; SLEB2; hPD-1; hPD-l; hSLE1   |
| <b>Dilution</b>    | WB~~ 1/500 - 1/2000 IHC~~1:100~500 ICC~~N/A E~~ 1/10000  |
| <b>Storage</b>     | Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| <b>Precautions</b> | PD1 is for research use only and not for use in diagnostic or therapeutic procedures.  |

## Protein Information

---

|                 |  |
|-----------------|--|
| <b>Name</b>     | PDCD1 {ECO:0000303 PubMed:7851902, ECO:0000312 HGNC:HGNC:8760}   |
| <b>Function</b> | Inhibitory receptor on antigen activated T-cells that plays a critical role in induction and maintenance of immune tolerance to self (PubMed: <a href="#">21276005</a> , PubMed: <a href="#">31754127</a> , PubMed: <a href="#">32184441</a> , PubMed: <a href="#">37208329</a> ). Delivers inhibitory signals upon binding to ligands CD274/PDCD1L1 and CD273/PDCD1LG2 (PubMed: <a href="#">21276005</a> , PubMed: <a href="#">26602187</a> ). Following T-cell receptor (TCR) engagement, PDCD1 associates with TCR-CD3 in the immunological synapse and directly inhibits T-cell activation (PubMed: <a href="#">32184441</a> ). Suppresses T-cell activation through the recruitment of PTPN11/SHP-2: following ligand-binding, PDCD1 is phosphorylated within the |

ITSM motif, leading to the recruitment of the protein tyrosine phosphatase PTPN11/SHP-2 that mediates dephosphorylation of key TCR proximal signaling molecules, such as ZAP70, PRKCQ/PKCtheta and CD247/CD3zeta (PubMed:[32184441](https://pubmed.ncbi.nlm.nih.gov/32184441/)).

## Cellular Location

Cell membrane; Single-pass type I membrane protein

## References

1. PLoS One. 2016 Jul 26;11(7):e0159383. 2. Tumour Biol. 2016 Jun;37(6):7507-14.

## Images




Figure 1: Black line: Control Antigen (100 ng); Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)




Figure 2: Western blot analysis using PD1 mAb against human PD1 (AA: extra 21-170) recombinant protein. (Expected MW is 46.6 kDa)





Figure 3: Western blot analysis using PD1 mouse mAb against MOLT4 (1), K562 (2), Ramos (3), HL-60 (4), THP-1 (5), and U937 (6) cell lysate.

Figure 4: Flow cytometric analysis of HeLa cells using PD1 mouse mAb (green) and negative control (red).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.