

Lipoprotein lipase Recombinant Rabbit mAb

Catalog # AP100011

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	P06858
Other Accession	P06858
Host	Rabbit
Clonality	Recombinant
Calculated MW	53162
Physical State	Liquid
Immunogen	A synthesized peptide derived from human Lipoprotein lipase
Epitope Specificity	445-475
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	10mM phosphate buffered saline(pH 7.4) with 150mM sodium chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol.
SIMILARITY	Belongs to the AB hydrolase superfamily. Lipase family.Contains 1 PLAT domain.
SUBUNIT	Homodimer. Interacts with APOC2; the interaction activates LPL activity in the presence of lipids. Interacts with GPIHBP1.
Post-translational modifications	Tyrosine nitration after lipopolysaccharide (LPS) challenge down-regulates the lipase activity.
DISEASE	Defects in LPL are the cause of lipoprotein lipase deficiency (LPL deficiency) [MIM:238600]; also known as familial chylomicronemia or hyperlipoproteinemia type I. LPL deficiency chylomicronemia is a recessive disorder usually manifesting in childhood. On a normal diet, patients often present with abdominal pain, hepatosplenomegaly, lipemia retinalis, eruptive xanthomata, and massive hypertriglyceridemia, sometimes complicated with acute pancreatitis.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Key enzyme in triglyceride metabolism. Catalyzes the hydrolysis of triglycerides from circulating chylomicrons and very low density lipoproteins (VLDL), and thereby plays an important role in lipid clearance from the blood stream, lipid utilization and storage.

Additional Information

Gene ID	4023
Other Names	HDLCQ11; LIPD; LIPL_BOVIN; LPL; Phospholipase A1; 3.1.1.34; LIPL_HUMAN; LIPL_MOUSE; lipoprotein lipase
Dilution	WB=1:1000-2000,IHC-P=1:100-200,IHC-F=1:100-200,IF=1:100-200
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	LPL
Synonyms	LIPD
Function	Key enzyme in triglyceride metabolism. Catalyzes the hydrolysis of triglycerides from circulating chylomicrons and very low density lipoproteins (VLDL), and thereby plays an important role in lipid clearance from the blood stream, lipid utilization and storage (PubMed:11342582, PubMed:27578112, PubMed:8675619). Although it has both phospholipase and triglyceride lipase activities it is primarily a triglyceride lipase with low but detectable phospholipase activity (PubMed:12032167, PubMed:7592706). Mediates margination of triglyceride-rich lipoprotein particles in capillaries (PubMed:24726386). Recruited to its site of action on the luminal surface of vascular endothelium by binding to GPIHBP1 and cell surface heparan sulfate proteoglycans (PubMed:11342582, PubMed:27811232).
Cellular Location	Cell membrane {ECO:0000250 UniProtKB:P11151}; Peripheral membrane protein {ECO:0000250 UniProtKB:P11151}; Extracellular side {ECO:0000250 UniProtKB:P11151}. Secreted. Secreted, extracellular space, extracellular matrix. Note=Newly synthesized LPL binds to cell surface heparan proteoglycans and is then released by heparanase Subsequently, it becomes attached to heparan proteoglycan on endothelial cells (PubMed:27811232). Locates to the plasma membrane of microvilli of hepatocytes with triglyceride-rich lipoproteins (TRL) Some of the bound LPL is then internalized and located inside non- coated endocytic vesicles (By similarity) {ECO:0000250 UniProtKB:P11151, ECO:0000269 PubMed:27811232}
Tissue Location	Detected in blood plasma (PubMed:11893776, PubMed:12641539, PubMed:2340307). Detected in milk (at protein level) (PubMed:2340307).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.