

WDR5 Recombinant Rabbit mAb

Catalog # AP100021

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	P61964
Other Accession	P61964
Host	Rabbit
Clonality	Recombinant
Calculated MW	36588
Physical State	Liquid
Immunogen	A synthesized peptide derived from human WDR5
Epitope Specificity	302-334/334
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	10mM phosphate buffered saline(pH 7.4) with 150mM sodium chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Contributes to histone modification. May position the N-terminus of histone H3 for efficient trimethylation at 'Lys-4'. As part of the MLL1/MLL complex it is involved in methylation and dimethylation at 'Lys-4' of histone H3. H3 'Lys-4' methylation represents a specific tag for epigenetic transcriptional activation.

Additional Information

Gene ID	11091
Other Names	WDR5_HUMAN; WDR5; BMP2-induced 3-kb gene protein; BIG3; BIG-3; CFAP89; SWD3; WD repeat domain 5; SWD3, Set1c WD40 repeat protein, homolog (S. cerevisiae); cilia and flagella associated protein 89; BMP-2-induced gene 3 kb
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,Flow-Cyt=1:50-100,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	WDR5
Synonyms	BIG3

Function	Contributes to histone modification (PubMed: 16600877 , PubMed: 16829960 , PubMed: 19103755 , PubMed: 19131338 , PubMed: 19556245 , PubMed: 20018852). May position the N-terminus of histone H3 for efficient trimethylation at 'Lys-4' (PubMed: 16829960). As part of the MLL1/MLL complex it is involved in methylation and dimethylation at 'Lys-4' of histone H3 (PubMed: 19556245). H3 'Lys-4' methylation represents a specific tag for epigenetic transcriptional activation (PubMed: 18840606). As part of the NSL complex it may be involved in acetylation of nucleosomal histone H4 on several lysine residues (PubMed: 19103755 , PubMed: 20018852). May regulate osteoblasts differentiation (By similarity). In association with RBBP5 and ASH2L, stimulates the histone methyltransferase activities of KMT2A, KMT2B, KMT2C, KMT2D, SETD1A and SETD1B (PubMed: 21220120 , PubMed: 22266653).
Cellular Location	Nucleus

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.