

VPS24 Recombinant Rabbit mAb

Catalog # AP100030

Product Information

Application	WB, IHC-P, IHC-F, IF, IP
Primary Accession	Q9Y3E7
Other Accession	Q9Y3E7
Host	Rabbit
Clonality	Recombinant
Calculated MW	25073
Physical State	Liquid
Immunogen	A synthesized peptide derived from human VPS24
Epitope Specificity	60-222/222
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	10mM phosphate buffered saline(pH 7.4) with 150mM sodium chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol.
SIMILARITY	Belongs to the SNF7 family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Probable core component of the endosomal sorting required for transport complex III (ESCRT-III) which is involved in multivesicular bodies (MVBs) formation and sorting of endosomal cargo proteins into MVBs. MVBs contain intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids.

Additional Information

Gene ID	51652
Other Names	VPS24; CGI-149; NEDF; 25.1; 4921505F14Rik; 9130011K15Rik; CGI-49; D6Ert286e; fb47g01; wu:fb47g01; zgc:76972; CHMP3_BOVIN; CHMP3; Chromatin-modifying protein 3; Vacuolar protein sorting-associated protein 24; CHMP3_HUMAN; Neuroendocrine differentiation factor; Vacuolar protein sorting-associated protein 24 (hVps24); CGI149; CHMP3_MOUSE; CHMP3_RAT; Vacuolar protein sorting-associated protein 24 (rVps24p); CHMP3_DANRE; Vacuolar protein-sorting-associated protein 24; charged multivesicular body protein 3; vacuolar protein sorting 24 (yeast); vacuolar protein sorting 24 homolog (S. cerevisiae)
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IP=1:20-50,Flow-Cyt=1:50-100,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name	CHMP3
Synonyms	CGI149, NEDF, VPS24
Function	Probable core component of the endosomal sorting required for transport complex III (ESCRT-III) which is involved in multivesicular bodies (MVBs) formation and sorting of endosomal cargo proteins into MVBs. MVBs contain intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids. The MVB pathway appears to require the sequential function of ESCRT-O, -I, -II and -III complexes. ESCRT-III proteins mostly dissociate from the invaginating membrane before the ILV is released. The ESCRT machinery also functions in topologically equivalent membrane fission events, such as the terminal stages of cytokinesis and the budding of enveloped viruses (HIV-1 and other lentiviruses). ESCRT-III proteins are believed to mediate the necessary vesicle extrusion and/or membrane fission activities, possibly in conjunction with the AAA ATPase VPS4. Selectively binds to phosphatidylinositol 3,5-bisphosphate PtdIns(3,5)P₂ and PtdIns(3,4)P₂ in preference to other phosphoinositides tested. Involved in late stages of cytokinesis. Plays a role in endosomal sorting/trafficking of EGF receptor. Isoform 2 prevents stress-mediated cell death and accumulation of reactive oxygen species when expressed in yeast cells.
Cellular Location	Cytoplasm, cytosol. Membrane; Lipid-anchor. Endosome. Late endosome membrane. Note=Localizes to the midbody of dividing cells
Tissue Location	Widely expressed. Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.