10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ZNF259/ZPR1 Recombinant Rabbit mAb

Catalog # AP100047

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	O75312
Other Accession	O75312
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Recombinant
Calculated MW	50925
Physical State	Liquid
Immunogen	A synthesized peptide derived from human ZPR1
Epitope Specificity	400-459
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	10mM phosphate buffered saline(pH 7.4) with 150mM sodium chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol.
SIMILARITY	Belongs to the ZPR1 family.
SUBUNIT	Component of an import snRNP complex composed of KPNB1, SNUPN, SMN1 and ZNF259. Interacts (via C-terminal region) with SMN1 (via C-terminal region); the interaction occurs after treatment with serum By similarity. Interacts with elongation factor 1-alpha EEF1A1; the interaction occurs in a epidermal growth factor (EGF)-dependent manner. Interacts (via zinc fingers) with EGFR (via C-terminal cytoplasmic kinase domain); the interaction is negatively regulated in response to epidermal growth factor (EGF) stimulation and EGFR kinase activity. May also bind to the PDGFR receptor.
DISEASE	May contribute to the severity of spinal muscular atrophy by increasing spinal motor neurons degeneration (Ref.7). Disruption phenotype Die during early embryonic development. Embryos show growth delay, failed to form normal trophectoderm and to expand the inner cell mass.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Plays a role for the localization and accumulation of the survival motor neuron protein SMN1 in sub-nuclear bodies, including gems and Cajal bodies. Induces neuron differentiation and stimulates axonal growth and formation of growth cone in spinal cord motor neurons. Plays a role in the splicing of cellular pre-mRNAs. May be involved in H2O2-induced neuronal cell death.

Additional Information

Gene ID	8882
Other Names	GKAF; ZNF259; Zfp259; RGD1562173; ZPR1_HUMAN; ZPR1; Zinc finger protein 259; ZPR1_MOUSE; ZPR1 zinc finger
Dilution	WB=1:1000-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500

Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	ZPR1
Synonyms	ZNF259
Function	Acts as a signaling molecule that communicates proliferative growth signals from the cytoplasm to the nucleus. It is involved in the positive regulation of cell cycle progression (PubMed:29851065). Plays a role for the localization and accumulation of the survival motor neuron protein SMN1 in sub-nuclear bodies, including gems and Cajal bodies. Induces neuron differentiation and stimulates axonal growth and formation of growth cone in spinal cord motor neurons. Plays a role in the splicing of cellular pre-mRNAs. May be involved in H(2)O(2)-induced neuronal cell death.
Cellular Location	Nucleus. Nucleus, nucleolus. Nucleus, gem. Nucleus, Cajal body. Cytoplasm, perinuclear region. Cytoplasm. Cell projection, axon. Cell projection, growth cone. Note=Colocalized with SMN1 in Gemini of coiled bodies (gems), Cajal bodies, axon and growth cones of neurons (By similarity) Localized predominantly in the cytoplasm in serum-starved cells growth arrested in G0 of the mitotic cell cycle. Localized both in the nucleus and cytoplasm at the G1 phase of the mitotic cell cycle. Accumulates in the subnuclear bodies during progression into the S phase of the mitotic cell cycle. Diffusely localized throughout the cell during mitosis. Colocalized with NPAT and SMN1 in nuclear bodies including gems (Gemini of coiled bodies) and Cajal bodies in a cell cycle- dependent manner. Translocates together with EEF1A1 from the cytoplasm to the nucleolus after treatment with mitogens. Colocalized with EGFR in the cytoplasm of quiescent cells. Translocates from the cytoplasm to the nucleus in a epidermal growth factor (EGF)-dependent manner
Tissue Location	Expressed in fibroblast; weakly expressed in fibroblast of spinal muscular atrophy (SMA) patients

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.