

IGJ Recombinant Rabbit mAb

Catalog # AP100081

Product Information

Application	WB, IHC-P, IHC-F, IF, IP, E
Primary Accession	P01591
Other Accession	P01591
Host	Rabbit
Clonality	Recombinant
Calculated MW	18099
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBUNIT	Secreted.
Background Descriptions	Enables IgA binding activity and protein homodimerization activity. Contributes to several functions, including immunoglobulin receptor binding activity; peptidoglycan binding activity; and phosphatidylcholine binding activity. Involved in several processes, including defense response to other organism; glomerular filtration; and positive regulation of respiratory burst. Located in extracellular space. Part of monomeric IgA immunoglobulin complex; pentameric IgM immunoglobulin complex; and secretory dimeric IgA immunoglobulin complex. [provided by Alliance of Genome Resources, Nov 2021]

Additional Information

Gene ID	3512
Other Names	IGCJ; IGJ; JCH; IGJ_HUMAN; JCHAIN; Joining chain of multimeric IgA and IgM; immunoglobulin J polypeptide, linker protein for immunoglobulin alpha and mu polypeptides; immunoglobulin J chain; IgJ chain; J chain
Dilution	IHC-P=1:50-200,IHC-F=1:50-200,WB=1:500-2000,IF=1:50-200,ELISA=1:5000-10 000,IP=1:50-200
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	JCHAIN (HGNC:5713)
Synonyms	IGCJ, IGJ

Function	Serves to link two monomer units of either IgM or IgA. In the case of IgM, the J chain-joined dimer is a nucleating unit for the IgM pentamer, and in the case of IgA it induces dimers and/or larger polymers. It also helps to bind these immunoglobulins to secretory component.
Cellular Location	Secreted {ECO:0000250 UniProtKB:P01592}.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.