

BSND Recombinant Rabbit mAb

Catalog # AP100089

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q8WZ55
Other Accession	Q8WZ55
Host	Rabbit
Clonality	Recombinant
Calculated MW	35197
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Background Descriptions	This gene encodes an essential beta subunit for CLC chloride channels. These heteromeric channels localize to basolateral membranes of renal tubules and of potassium-secreting epithelia of the inner ear. Mutations in this gene have been associated with Bartter syndrome with sensorineural deafness. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID	7809
Other Names	Barttin;BART
Dilution	IHC-P=1:50-200,IHC-F=1:50-200,WB=1:500-2000,IF=1:50-200
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	BSND {ECO:0000303 PubMed:11687798, ECO:0000312 HGNC:HGNC:16512}
Function	Regulatory subunit of anion-selective CLCNKA:BSND and CLCNKB:BSND heteromeric channels involved in basolateral chloride conductance along the nephron to achieve urine concentration and maintain systemic acid-base homeostasis, and in the stria vascularis of the inner ear to establish the endocochlear potential necessary for normal hearing (PubMed: 11734858 , PubMed: 12111250 , PubMed: 12574213 , PubMed: 16849430 , PubMed: 18776122 , PubMed: 19646679 , PubMed: 20538786 , PubMed: 26013830). Most likely acts as a chaperone that allosterically regulates proper sorting of CLCNKA:BSND and CLCNKB:BSND channels at the basolateral plasma membrane domain and functional switch to ion

conducting state. Mediates constitutive opening of channel common gates (PubMed:[11734858](#), PubMed:[12111250](#), PubMed:[12574213](#), PubMed:[16849430](#), PubMed:[18776122](#), PubMed:[19646679](#), PubMed:[20538786](#), PubMed:[26013830](#)).

Cellular Location

Basolateral cell membrane; Multi-pass membrane protein. Note=Mostly sorted at the basolateral membrane. A significant amount also observed intracellularly. Staining in membranes of the renal tubule and of potassium-secreting epithelia of the inner ear is basolateral (By similarity).

Tissue Location

Expressed primarily in kidney. Expressed in specific nephron segments and in the stria vascularis of the inner ear

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.