

DDX4 Recombinant Rabbit mAb

Catalog # AP100100

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9NQI0
Other Accession	Q9NQI0
Host	Rabbit
Clonality	Recombinant
Calculated MW	79308
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human DDX4
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Additional Information

Gene ID	54514
Other Names	DDX 4; DEAD(Asp Glu Ala Asp) box polypeptide 4; Dead box protein 4; Dead-box protein 4; MGC111074; MVH; Probable ATP dependent RNA helicase DDX4; VASA; VASA homolog; DDX4_HUMAN.; DDX4;
Dilution	IHC-P=1:200-800,IHC-F=1:200-800,IF=1:200-800
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	DDX4
Synonyms	VASA
Function	ATP-dependent RNA helicase required during spermatogenesis (PubMed: 10920202 , PubMed: 21034600). Required to repress transposable elements and preventing their mobilization, which is essential for the germline integrity (By similarity). Acts via the piRNA metabolic process, which mediates the repression of transposable elements during meiosis by forming complexes composed of piRNAs and Piwi proteins and governs the methylation and subsequent repression of transposons (By similarity). Involved in the secondary piRNAs metabolic process, the production of piRNAs in fetal male germ cells through a ping-pong amplification cycle (By similarity). Required for PIWIL2 slicing- triggered piRNA biogenesis: helicase

activity enables utilization of one of the slice cleavage fragments generated by PIWIL2 and processing these pre-piRNAs into piRNAs (By similarity).

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q61496}. Cytoplasm, perinuclear region {ECO:0000250|UniProtKB:Q61496} Note=Component of the meiotic nuage, also named P granule, a germ-cell- specific organelle required to repress transposon activity during meiosis. {ECO:0000250|UniProtKB:Q61496}

Tissue Location

Expressed only in ovary and testis. Expressed in migratory primordial germ cells in the region of the gonadal ridge in both sexes.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.