

Histone H1oo Rabbit pAb

Catalog # AP100150

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q8IZA3
Other Accession	Q8IZA3
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35813
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human H1FOO
Epitope Specificity	81-180/346
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the histone H1/H5 family. Contains 1 H15 (linker histone H1/H5 globular) domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. The protein encoded is a member of the histone H1 family. This gene contains introns, unlike most histone genes. The protein encoded is a member of the histone H1 family. The related mouse gene is expressed only in oocytes. [provided by RefSeq, Jul 2008].

Additional Information

Gene ID	132243
Other Names	H1.8; H1FOO; H1oo; osH1; H18_HUMAN; H1-8; Histone H1oo; Oocyte-specific histone H1; Oocyte-specific linker histone H1 (osH1); H1.8 linker histone; H1 histone family, member O, oocyte-specific; H1 histone family member O oocyte specific
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	H1-8 (HGNC:18463)
Function	May play a key role in the control of gene expression during oogenesis and early embryogenesis, presumably through the perturbation of chromatin structure. Essential for meiotic maturation of germinal vesicle-stage oocytes. The somatic type linker histone H1c is rapidly replaced by H1oo in a donor nucleus transplanted into an oocyte. The greater mobility of H1oo as compared to H1c may contribute to this rapid replacement and increased instability of the embryonic chromatin structure. The rapid replacement of H1c with H1oo may play an important role in nuclear remodeling (By similarity).
Cellular Location	Cytoplasm. Nucleus {ECO:0000255 PROSITE-ProRule:PRU00837}. Chromosome {ECO:0000255 PROSITE-ProRule:PRU00837}
Tissue Location	Oocyte-specific..

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.