

CEP152 Rabbit pAb

Catalog # AP100211

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	O94986
Other Accession	O94986
Host	Rabbit
Clonality	Polyclonal
Calculated MW	195626
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human CEP152
Epitope Specificity	901-1000/1654
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBUNIT	Interacts (via N-terminus) with PLK4. Interacts (via C-terminus) with CENPJ (via-N-terminus). Interacts with CINP. Interacts with CEP63; this interaction recruits CEP152 to centrosomes.
DISEASE	Defects in CEP152 are the cause of microcephaly primary type 4 (MCPH4) [MIM:604321]. A disease defined as a head circumference more than 3 standard deviations below the age-related mean. Brain weight is markedly reduced and the cerebral cortex is disproportionately small. Despite this marked reduction in size, the gyral pattern is relatively well preserved, with no major abnormality in cortical architecture. Affected individuals are mentally retarded. Primary microcephaly is further defined by the absence of other syndromic features or significant neurological deficits due to degenerative brain disorder. Defects in CEP152 are the cause of Seckel syndrome type 5 (SCKL5) [MIM:613823]. A rare autosomal recessive disorder characterized by proportionate dwarfism of prenatal onset associated with low birth weight, growth retardation, severe microcephaly with a bird-headed like appearance, and mental retardation.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Defects in CEP152 are the cause of microcephaly primary type 4 (MCPH4). A disease defined as a head circumference more than 3 standard deviations below the age-related mean. Brain weight is markedly reduced and the cerebral cortex is disproportionately small. Despite this marked reduction in size, the gyral pattern is relatively well preserved, with no major abnormality in cortical architecture. Affected individuals are mentally retarded. Primary microcephaly is further defined by the absence of other syndromic features or significant neurological deficits due to degenerative brain disorder.

Additional Information

Gene ID 22995

Other Names	MCPH4; MCPH9; SCKL5; mKIAA0912; CE152_HUMAN; CEP152; KIAA0912; CE152_MOUSE; centrosomal protein 152; centrosomal protein 152kDa; asterless
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	CEP152 (HGNC:29298)
Synonyms	KIAA0912
Function	Necessary for centrosome duplication; the function also seems to involve CEP63, CDK5RAP2 and WDR62 through a stepwise assembled complex at the centrosome that recruits CDK2 required for centriole duplication (PubMed: 26297806). Acts as a molecular scaffold facilitating the interaction of PLK4 and CPAP, 2 molecules involved in centriole formation (PubMed: 20852615 , PubMed: 21059844). Proposed to snatch PLK4 away from PLK4:CEP92 complexes in early G1 daughter centriole and to reposition PLK4 at the outer boundary of a newly forming CEP152 ring structure (PubMed: 24997597). Also plays a key role in deuterosome-mediated centriole amplification in multiciliated that can generate more than 100 centrioles (By similarity). Overexpression of CEP152 can drive amplification of centrioles (PubMed: 20852615).
Cellular Location	Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole Note=Colocalizes with CDK5RAP2, WDR62 and CEP63 in a discrete ring around the proximal end of the parental centriole. At this site, a cohesive structure is predicted to engage parental centrioles and procentrioles (PubMed:21983783, PubMed:26297806). Localizes to the deuterosome (By similarity). Localizes to pericentriolar material (PCM) (PubMed:26337392). {ECO:0000250 UniProtKB:Q498G2, ECO:0000269 PubMed:21983783, ECO:0000269 PubMed:26297806, ECO:0000269 PubMed:26337392}

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.