

VHR Recombinant Rabbit mAb

Catalog # AP100299

Product Information

Application	WB, IHC-P, IHC-F, IF, IP, E
Primary Accession	P51452
Other Accession	P51452
Host	Rabbit
Clonality	Recombinant
Calculated MW	20478
Isotype	IgG
Purity	affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SIMILARITY Belongs to the protein-tyrosine phosphatase family. Non-receptor class dual specificity subfamily. Contains 1 tyrosine-protein phosphatase domain.

SUBUNIT Interacts with VRK3, which seems to activate its phosphatase activity

Background Descriptions The protein encoded by this gene is a member of the dual specificity protein phosphatase subfamily. These phosphatases inactivate their target kinases by dephosphorylating both the phosphoserine/threonine and phosphotyrosine residues. They negatively regulate members of the mitogen-activated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which are associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate specificities for various MAP kinases, different tissue distribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli. This gene maps in a region that contains the BRCA1 locus which confers susceptibility to breast and ovarian cancer. Although DUSP3 is expressed in both breast and ovarian tissues, mutation screening in breast ca

Additional Information

Gene ID	1845
Other Names	VHR; 2210015O03Rik; 5031436O03Rik; T-DSP11; RGD1560049; DUS3_HUMAN; DUSP3; Dual specificity protein phosphatase VHR; Vaccinia H1-related phosphatase (VHR); 3.1.3.16; 3.1.3.48; DUS3_MOUSE; dual specificity phosphatase 3; vaccinia virus phosphatase VH1-related; Dual specificity protein phosphatase 3; Vaccinia H1-related phosphatase
Dilution	IHC-P=1:50-200, IHC-F=1:50-200, WB=1:500-2000, IF=1:50-200, ELISA=1:5000-10000, IP=1:50-200
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	DUSP3
Synonyms	VHR
Function	Shows activity both for tyrosine-protein phosphate and serine-protein phosphate, but displays a strong preference toward phosphotyrosines (PubMed: 10224087 , PubMed: 11863439). Specifically dephosphorylates and inactivates ERK1 and ERK2 (PubMed: 10224087 , PubMed: 11863439).
Cellular Location	Nucleus. Cytoplasm, cytoskeleton, flagellum axoneme {ECO:0000250 UniProtKB:Q9D7X3}

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.