

GAD-65/67 Recombinant Rabbit mAb

Catalog # AP100302

Product Information

Application	WB, IHC-P, IHC-F, IF, IP, E
Primary Accession	Q05329
Other Accession	Q05329
Host	Rabbit
Clonality	Recombinant
Calculated MW	65411
Isotype	IgG
Purity	affinity purified by Protein A

Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Background Descriptions	glutamate decarboxylase 1(GAD1) Homo sapiens This gene encodes one of several forms of glutamic acid decarboxylase, identified as a major autoantigen in insulin-dependent diabetes. The enzyme encoded is responsible for catalyzing the production of gamma-aminobutyric acid from L-glutamic acid. A pathogenic role for this enzyme has been identified in the human pancreas since it has been identified as an autoantigen and an autoreactive T cell target in insulin-dependent diabetes. This gene may also play a role in the stiff man syndrome. Deficiency in this enzyme has been shown to lead to pyridoxine dependency with seizures. Alternative splicing of this gene results in two products, the predominant 67-kD form and a less-frequent 25-kD form. [provided by RefSeq, Jul 2008],

Additional Information

Gene ID	2572
Other Names	Glutamate decarboxylase 1/2;GAD1;GAD;GAD67;Glutamate decarboxylase 1;67 kDa glutamic acid decarboxylase;GAD-67;Glutamate decarboxylase 67 kDa isoform;GAD2;GAD65;Glutamate decarboxylase 2;65 kDa glutamic acid decarboxylase;GAD-65;Glutamate decarboxylase 65
Dilution	IHC-P=1:50-200,IHC-F=1:50-200,WB=1:500-2000,IF=1:50-200,ELISA=1:5000-10000,IP=1:50-200
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	GAD2 (HGNC:4093)
-------------	------------------------------------

Synonyms	GAD65
Function	Catalyzes the production of GABA.
Cellular Location	Cytoplasm, cytosol. Cytoplasmic vesicle. Presynaptic cell membrane; Lipid-anchor. Golgi apparatus membrane; Peripheral membrane protein; Cytoplasmic side. Note=Associated to cytoplasmic vesicles In neurons, cytosolic leaflet of Golgi membranes and presynaptic clusters

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.