

VE-Cadherin Recombinant Rabbit mAb

Catalog # AP100321

Product Information

Application	WB, IP, E
Primary Accession	P33151
Other Accession	P33151
Host	Rabbit
Clonality	Recombinant
Calculated MW	87528
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 5 cadherin domains.
SUBUNIT	Interacts via cadherin 5 domain with PTPRB. Interacts with TRPC4. Interacts with KRIT1.
Post-translational modifications	Phosphorylated on tyrosine residues by KDR/VEGFR-2. Dephosphorylated by PTPRB.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a classical cadherin of the cadherin superfamily. The encoded preproprotein is proteolytically processed to generate the mature glycoprotein. This calcium-dependent cell-cell adhesion molecule is comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. Functioning as a classical cadherin by imparting to cells the ability to adhere in a homophilic manner, this protein plays a role in endothelial adherens junction assembly and maintenance. This gene is located in a gene cluster in a region on the long arm of chromosome 16 that is involved in loss of heterozygosity events in breast and prostate cancer. [provided by RefSeq, Nov 2015],

Additional Information

Gene ID	1003
Other Names	7B4; CD144; VE-Cad; VECD; VEcad; Vec; CADH5_HUMAN; CDH5; 7B4 antigen; Vascular endothelial cadherin (VE-cadherin); CADH5_MOUSE; cadherin 5; cadherin 5, type 2, VE-cadherin (vascular epithelium); cadherin 5, type 2 (vascular endothelium); VE-cadherin; Cadherin-5; Vascular endothelial cadherin; CD antigen CD144
Dilution	WB=1:500-2000,ELISA=1:5000-10000,IP=1:50-200
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	CDH5 (HGNC:1764)
Function	Cadherins are calcium-dependent cell adhesion proteins (By similarity). They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types (PubMed:21269602). This cadherin may play a important role in endothelial cell biology through control of the cohesion and organization of the intercellular junctions (By similarity). It associates with alpha-catenin forming a link to the cytoskeleton (PubMed:10861224). Plays a role in coupling actin fibers to cell junctions in endothelial cells, via acting as a cell junctional complex anchor for AMOTL2 and MAGI1 (By similarity). Acts in concert with KRIT1 and PALS1 to establish and maintain correct endothelial cell polarity and vascular lumen (By similarity). These effects are mediated by recruitment and activation of the Par polarity complex and RAP1B (PubMed:20332120). Positively regulates reorientation of actin stress fibers and endothelial cell reorientation in response to cellular mechanotransduction (PubMed:25795300). Required for activation of PRKCZ and for the localization of phosphorylated PRKCZ, PARD3, TIAM1 and RAP1B to the cell junction (PubMed:20332120). Associates with CTNND1/p120-catenin to control CADH5 endocytosis (By similarity).
Cellular Location	Cell junction, adherens junction. Cell membrane; Single-pass type I membrane protein Cytoplasm {ECO:0000250 UniProtKB:P55284}. Note=Found at cell-cell boundaries and probably at cell-matrix boundaries. KRIT1 and CDH5 reciprocally regulate their localization to endothelial cell-cell junctions.
Tissue Location	Expressed in endothelial cells (at protein level) (PubMed:27338829). Expressed in the brain (PubMed:2059658)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.