10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-CD8-AcalephFluor488-labled

Catalog # AP100364

Product Information

Application	FC, IF/IC
Primary Accession	P01731, P10300
Reactivity	Mouse
Clonality	Monoclonal
Calculated MW	27456

Additional Information

Gene ID	12525
Other Names	MAL; T-cell surface glycoprotein CD8 alpha chain; T-lymphocyte differentiation antigen T8/Leu-2; CD antigen CD8a; CD8B1; T-cell surface glycoprotein CD8 beta chain; CD antigen CD8b
Target/Specificity	Recognizes mouse CD8
Dilution	FC~~1:10~50 IF/IC~~N/A
Format	Rat IgG2b kappa. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	Cd8a
Synonyms	Lyt-2, Lyt2
Function	Integral membrane glycoprotein that plays an essential role in the immune response and serves multiple functions in responses against both external and internal offenses. In T-cells, functions primarily as a coreceptor for MHC class I molecule:peptide complex. The antigens presented by class I peptides are derived from cytosolic proteins while class II derived from extracellular proteins. Interacts simultaneously with the T-cell receptor (TCR) and the MHC class I proteins presented by antigen presenting cells (APCs). In turn, recruits the Src kinase LCK to the vicinity of the TCR-CD3 complex. LCK then initiates different intracellular signaling pathways by phosphorylating various substrates ultimately leading to lymphokine production, motility, adhesion and activation of cytotoxic T- lymphocytes (CTLs). This mechanism enables CTLs to recognize and eliminate infected cells and tumor cells. In NK-cells, the presence of CD8A homodimers at the cell surface provides a survival mechanism allowing conjugation and lysis of multiple target cells. CD8A homodimer molecules also promote the survival and differentiation of

	activated lymphocytes into memory CD8 T-cells.
Cellular Location	Cell membrane {ECO:0000250|UniProtKB:P01732}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:P01732} Note=Cd8a localizes to lipid rafts only when associated with its partner Cd8b. {ECO:0000250|UniProtKB:P01732}

Background

AcalephFluor488-labled rat monoclonal antibody to CD8

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.