10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Anti-CD8-FITC-labled

Catalog # AP100370

## Product Information

| | |
|---|---|
| **Application** | FC, IF/IC |
| **Primary Accession** | P01731, P10300 |
| **Reactivity** | Mouse |
| **Clonality** | Monoclonal |
| **Calculated MW** | 27456 |

## Additional Information

| | |
|---|---|
| **Gene ID** | 12525 |
| **Other Names** | MAL; T-cell surface glycoprotein CD8 alpha chain; T-lymphocyte differentiation antigen T8/Leu-2; CD antigen CD8a; CD8B1; T-cell surface glycoprotein CD8 beta chain; CD antigen CD8b |
| **Target/Specificity** | Recognizes mouse CD8 |
| **Dilution** | FC~~1:10~50 IF/IC~~N/A |
| **Format** | Rat IgG2b kappa. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide. |
| **Storage** | Store at -20 °C.Stable for 12 months from date of receipt |

## Protein Information

| | |
|---|---|
| **Name** | Cd8a |
| **Synonyms** | Lyt-2, Lyt2 |
| **Function** | Integral membrane glycoprotein that plays an essential role in the immune response and serves multiple functions in responses against both external and internal offenses. In T-cells, functions primarily as a coreceptor for MHC class I molecule:peptide complex. The antigens presented by class I peptides are derived from cytosolic proteins while class II derived from extracellular proteins. Interacts simultaneously with the T-cell receptor (TCR) and the MHC class I proteins presented by antigen presenting cells (APCs). In turn, recruits the Src kinase LCK to the vicinity of the TCR-CD3 complex. LCK then initiates different intracellular signaling pathways by phosphorylating various substrates ultimately leading to lymphokine production, motility, adhesion and activation of cytotoxic T- lymphocytes (CTLs). This mechanism enables CTLs to recognize and eliminate infected cells and tumor cells. In NK-cells, the presence of CD8A homodimers at the cell surface provides a survival mechanism allowing conjugation and lysis of multiple target cells. CD8A homodimer molecules also promote the survival and differentiation of |

| | |
|---|---|
| | activated lymphocytes into memory CD8 T-cells. |
| **Cellular Location** | Cell membrane {ECO:0000250\|UniProtKB:P01732}; Single-pass type I membrane protein {ECO:0000250\|UniProtKB:P01732} Note=Cd8a localizes to lipid rafts only when associated with its partner Cd8b. {ECO:0000250\|UniProtKB:P01732} |

## Background

FITC-labled rat monoclonal antibody to CD8

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.