10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-CD18-AcalephFluor647-labled

Catalog # AP100528

Product Information

Application	FC, IF/IC
Primary Accession	P05107
Reactivity	Human
Clonality	Monoclonal
Calculated MW	84791

Additional Information

Gene ID	3689
Other Names	CD18; MFI7; Integrin beta-2; Cell surface adhesion glycoproteins LFA-1/CR3/p150.95 subunit beta; Complement receptor C3 subunit beta; CD antigen CD18
Target/Specificity	Recognizes human CD18
Dilution	FC~~1:10~50 IF/IC~~N/A
Format	Mouse IgG1. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	ITGB2 (HGNC:6155)
Synonyms	CD18, MFI7
Function	Integrin ITGAL:ITGB2 is a receptor for ICAM1, ICAM2 and ICAM3 (PubMed:1676048, PubMed:23775590, PubMed:38195629). Integrin ITGAL:ITGB2 is also a receptor for the secreted form of ubiquitin-like protein ISG15; the interaction is mediated by ITGAL (PubMed:29100055). Integrins ITGAM:ITGB2 and ITGAX:ITGB2 are receptors for the iC3b fragment of the third complement component and for fibrinogen. Integrin ITGAX:ITGB2 recognizes the sequence G-P-R in fibrinogen alpha-chain. Integrin ITGAM:ITGB2 recognizes P1 and P2 peptides of fibrinogen gamma chain. Integrin ITGAM:ITGB2 is also a receptor for factor X. Integrin ITGAD:ITGB2 is a receptor for ICAM3 and VCAM1 (PubMed:10438935, PubMed:8777714, PubMed:9841932). Contributes to natural killer cell cytotoxicity (PubMed:15356110). Involved in leukocyte adhesion and transmigration of leukocytes including T-cells and neutrophils (PubMed:11812992, PubMed:28807980). Triggers neutrophil transmigration during lung injury through PTK2B/PYK2-mediated activation (PubMed:18587400). Integrin

	ITGAL:ITGB2 in association with ICAM3, contributes to apoptotic neutrophil phagocytosis by macrophages (PubMed:23775590). In association with alpha subunit ITGAM/CD11b, required for CD177-PRTN3-mediated activation of TNF primed neutrophils (PubMed:21193407). Integrins ITGAX:ITGB2 functions as a receptor of the erythrocyte-specific adhesion molecule ICAM4 and mediates erythrophagocytosis (PubMed:16985175). Integrins ITGAX:ITGB2 functions as a receptor of the neuron-specific adhesion molecule ICAM5 ensuring neuron cell-leukocyte adhesion (PubMed:10741396). Integrin ITGAL:ITGB2 functions as a receptor of ICAM1 by acting as a platform at the immunological synapse to translate TCR engagement and density of the ITGAL ligand ICAM1 into graded adhesion (PubMed:38195629). Integrin ITGAM:ITGB2/MAC-1 complex functions as a signaling receptor for the ligand receptor ICAM1, ensuring adhesion between stimulated neutrophils and stimulated endothelial cells (PubMed:1980124). Integrin ITGAL/ITGB2 that functions as a signaling receptor of ICAM2, ensuring leukocyte cell-cell adhesion on resting cells (PubMed:1676048).
Cellular Location	Cell membrane; Single-pass type I membrane protein. Membrane raft; Single-pass type I membrane protein
Tissue Location	Leukocytes (PubMed:23775590). Expressed in neutrophils (at protein level) (PubMed:21193407, PubMed:28807980)

Background

AcalephFluor647-labled mouse monoclonal antibody to CD18

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.