10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-CD278-AcalephFluor647-labled

Catalog # AP100608

Product Information

Application	FC, IF/IC
Primary Accession	Q9WVS0
Reactivity	Mouse
Clonality	Monoclonal
Calculated MW	22709

Additional Information

Gene ID	54167
Other Names	AILIM; Inducible T-cell costimulator; Activation-inducible lymphocyte immunomediatory molecule; CD278
Target/Specificity	Recognizes mouse CD278
Dilution	FC~~1:10~50 IF/IC~~N/A
Format	Rat IgG2b kappa. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	Icos
Synonyms	Ailim
Function	Stimulatory receptor expressed in activated or antigen- experienced T-cells that plays an important role in the immune response (PubMed:11343123). Upon binding to its ligand ICOSL expressed on antigen presenting cells (APCs), delivers costimulatory signals that enhances all basic T-cell responses to a foreign antigen, namely proliferation, secretion of lymphokines including IL10, up-regulation of molecules that mediate cell-cell interaction, and effective help for antibody secretion by B-cells (PubMed:10657606). Also acts as a costimulatory receptor critical for the differentiation of T follicular regulatory cells upon immune challenges such as viral infection (PubMed:36754569). Mechanistically, potentiates TCR-induced calcium flux by augmenting PLCG1 activation and actin remodeling (PubMed:27693916). In addition, activates PI3K signaling pathways independently of calcium flux (PubMed:19915142, PubMed:27693916). Essential both for efficient interaction between T and B-cells and for normal antibody responses to T-cell dependent antigens. Prevents the apoptosis of pre-activated T-cells. Plays a critical role in CD40- mediated class switching of immunoglobin isotypes

	(PubMed:11343122).
Cellular Location	Cell membrane {ECO:0000250|UniProtKB:Q9Y6W8}; Single-pass type I membrane protein
Tissue Location	Expressed on activated T-cells and resting memory T-cells. High expression seen in the thymic medulla and in the germinal centers and T-cell zones of lymph nodes and Peyer patches. Expressed at low levels in the spleen.

Background

AcalephFluor647-labled rat monoclonal antibody to CD278

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.