10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-CD279-AcalephFluor594-labled

Catalog # AP100615

Product Information

Application	FC, IF/IC
Primary Accession	Q02242
Reactivity	Mouse
Clonality	Monoclonal
Calculated MW	31842

Additional Information

Gene ID	18566
Other Names	PD1; Programmed cell death protein 1; Protein PD-1; hPD-1; CD279
Target/Specificity	Recognizes mouse CD279
Dilution	FC~~1:10~50 IF/IC~~N/A
Format	Rat IgG2a kappa. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	Pdcd1 {ECO:0000312|MGI:MGI:104879}
Function	Inhibitory receptor on antigen activated T-cells that plays a critical role in induction and maintenance of immune tolerance to self (PubMed:10485649, PubMed:11209085, PubMed:11698646, PubMed:21300912). Delivers inhibitory signals upon binding to ligands, such as CD274/PDCD1L1 and CD273/PDCD1LG2 (PubMed:11015443, PubMed:11224527, PubMed:18287011, PubMed:18641123, PubMed:22641383). Following T-cell receptor (TCR) engagement, PDCD1 associates with TCR-CD3 in the immunological synapse and directly inhibits T-cell activation (PubMed:22641383). Suppresses T-cell activation through the recruitment of PTPN11/SHP-2: following ligand-binding, PDCD1 is phosphorylated within the ITSM motif, leading to the recruitment of the protein tyrosine phosphatase PTPN11/SHP-2 that mediates dephosphorylation of key TCR proximal signaling molecules, such as ZAP70, PRKCQ/PKCtheta and CD247/CD3zeta (PubMed:11698646, PubMed:22641383). The PDCD1-mediated inhibitory pathway is exploited by tumors to attenuate anti-tumor immunity and facilitate tumor survival (By similarity).
Cellular Location	Cell membrane; Single-pass type I membrane protein

Tissue Location	Thymus-specific..

Background

AcalephFluor594-labled rat monoclonal antibody to CD279

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.