10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-CD28-AcalephFluor594-labled

Catalog # AP101047

Product Information

Application	FC, IF/IC
Primary Accession	P31041
Reactivity	Mouse
Clonality	Monoclonal
Calculated MW	25243

Additional Information

Gene ID	12487
Other Names	T-cell-specific surface glycoprotein CD28; TP44; CD28
Target/Specificity	Recognizes mouse CD28
Dilution	FC~~1:10~50 IF/IC~~N/A
Format	Syrian Hamster IgG. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	Cd28
Function	Receptor that plays a role in T-cell activation, proliferation, survival and the maintenance of immune homeostasis (PubMed:7688139). Functions not only as an amplifier of TCR signals but delivers unique signals that control intracellular biochemical events that alter the gene expression program of T-cells. Stimulation upon engagement of its cognate ligands CD80 or CD86 increases proliferation and expression of various cytokines in particular IL2 production in both CD4(+) and CD8(+) T-cell subsets. Mechanistically, ligation induces recruitment of protein kinase C-theta/PRKCQ and GRB2 leading to NF-kappa-B activation via both PI3K/Akt-dependent and -independent pathways. In conjunction with TCR/CD3 ligation and CD40L costimulation, enhances the production of IL4 and IL10 in T-cells.
Cellular Location	Cell membrane; Single-pass type I membrane protein {ECO:0000250|UniProtKB:P10747}

Background

AcalephFluor594-labled syrian hamster monoclonal antibody to CD28

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.