10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-CD317-AcalephFluor568-labled

Catalog # AP101054

Product Information

Application	FC, IF/IC
Primary Accession	Q8R2Q8
Reactivity	Mouse
Clonality	Monoclonal
Calculated MW	19152

Additional Information

Gene ID	69550
Other Names	Bone marrow stromal antigen 2; BST-2; HM1.24 antigen; Tetherin; CD317
Target/Specificity	Recognizes mouse CD317
Dilution	FC~~1:10~50 IF/IC~~N/A
Format	Rat IgG2b kappa. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	Bst2
Function	IFN-induced antiviral host restriction factor which efficiently blocks the release of diverse mammalian enveloped viruses by directly tethering nascent virions to the membranes of infected cells. Acts as a direct physical tether, holding virions to the cell membrane and linking virions to each other. The tethered virions can be internalized by endocytosis and subsequently degraded or they can remain on the cell surface. In either case, their spread as cell-free virions is restricted. Its target viruses belong to diverse families, including retroviridae: human immunodeficiency virus type 1 (HIV-1), mouse mammary tumor virus (MMTV) and murine leukemia virus (MLV), filoviridae: ebola virus (EBOV), arenaviridae: lassa virus (LASV), and rhabdoviridae: vesicular stomatitis virus (VSV). Can inhibit cell surface proteolytic activity of MMP14 causing decreased activation of MMP15 which results in inhibition of cell growth and migration. Can stimulate signaling by LILRA4/ILT7 and consequently provide negative feedback to the production of IFN by plasmacytoid dendritic cells in response to viral infection. Plays a role in the organization of the subapical actin cytoskeleton in polarized epithelial cells.
Cellular Location	Golgi apparatus, trans-Golgi network. Cell membrane; Single-pass type II membrane protein. Cell membrane; Lipid-anchor, GPI-anchor. Late

	endosome. Membrane raft. Cytoplasm Apical cell membrane. Note=Shuttles between the cell membrane, where it is present predominantly in membrane/lipid rafts, and the trans-Golgi network. Forms a complex with MMP14 and localizes to the cytoplasm (By similarity).
Tissue Location	In naive mice, specifically expressed on type I interferon-producing cells (at protein level)

Background

AcalephFluor568-labled rat monoclonal antibody to CD317

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.