10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-CD86-AcalephFluor405-labled

Catalog # AP101171

Product Information

Application	FC, IF/IC
Primary Accession	P42082
Reactivity	Mouse
Clonality	Monoclonal
Calculated MW	34666

Additional Information

Gene ID	12524
Other Names	CD28LG2; T-lymphocyte activation antigen CD86; Activation B7-2 antigen; B70; BU63; CTLA-4 counter-receptor B7.2; FUN-1; CD86
Target/Specificity	Recognizes mouse CD86
Dilution	FC~~1:10~50 IF/IC~~N/A
Format	Rat IgG2a kappa. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	Cd86
Function	Costimulatory molecule that belongs to the immunoglobulin superfamily that plays an important role in T-lymphocyte activation (By similarity). Acts as the primary auxiliary signal augmenting the MHC/TCR signal in naive T-cells by acting as a ligand for the CD28 receptor which is constitutively expressed on the cell surface of T- cells (By similarity). May play a critical role in the early events of T-cell activation and costimulation of naive T-cells, such as deciding between immunity and anergy that is made by T-cells within 24 hours after activation (By similarity). Also involved in the regulation of B cells function, plays a role in regulating the level of IgG(1) produced (PubMed:23241883). Upon CD40 engagement, activates NF-kappa-B signaling pathway via phospholipase C and protein kinase C activation (PubMed:23241883). Also acts as an inhibitor of T-cell activation by acting as a ligand for CTLA4, a decoy receptor, thereby blocking CD28- mediated T-cell priming (PubMed:7504059, PubMed:7513726).
Cellular Location	Cell membrane {ECO:0000250|UniProtKB:P42081}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:P42081}

Tissue Location	Expressed on the surfaces of antigen-presenting cells.

Background

AcalephFluor405-labled rat monoclonal antibody to CD86

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.